



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 31/415, C07C 321/02, C07D 233/56</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 96/30015</b> <b>(43) International Publication Date:</b> 3 October 1996 (03.10.96)
<b>(21) International Application Number:</b> PCT/US96/03980 <b>(22) International Filing Date:</b> 25 March 1996 (25.03.96) <b>(30) Priority Data:</b> 08/413,137 29 March 1995 (29.03.95) US 08/412,830 29 March 1995 (29.03.95) US <b>(60) Parent Applications or Grants</b> <b>(63) Related by Continuation</b> US 08/412,830 (CON) Filed on 29 March 1995 (29.03.95) US 08/413,137 (CON) Filed on 29 March 1995 (29.03.95) <b>(71) Applicant (for all designated States except US):</b> MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> CICCARONE, Terrence, M. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). WILLIAMS, Theresa, M. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). MacTOUGH, Suzanne,	C. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). <b>(74) Common Representative:</b> MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). <b>(81) Designated States:</b> AL, AM, AU, AZ, BB, BG, BR, BY, CA, CN, CZ, EE, GE, HU, IS, JP, KG, KR, KZ, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>	
<b>(54) Title:</b> INHIBITORS OF FARNESYL-PROTEIN TRANSFERASE <b>(57) Abstract</b> <p>The present invention comprises peptidomimetic compounds which comprise a suitably substituted aminoalkylbenzamide moiety. The instant compounds inhibit the farnesyl protein transferase enzyme and the farnesylation of certain proteins. Furthermore, the instant farnesyl protein transferase inhibitors differ from those previously described as inhibitors of farnesyl-protein transferase in that they do not have a thiol moiety. The lack of the thiol offers unique advantages in terms of improved pharmacokinetic behavior in animals, prevention of thiol-dependent chemical reactions, such as rapid autoxidation and disulfide formation with endogenous thiols, and reduced systemic toxicity. Further contained in this invention are chemotherapeutic compositions containing these farnesyl transferase inhibitors and methods for their production.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic			SE	Sweden
CG	Congo	KR	Republic of Korea	SG	Singapore
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LR	Liberia	SZ	Swaziland
CS	Czechoslovakia	LT	Lithuania	TD	Chad
CZ	Czech Republic	LU	Luxembourg	TG	Togo
DE	Germany	LV	Latvia	TJ	Tajikistan
DK	Denmark	MC	Monaco	TT	Trinidad and Tobago
EE	Estonia	MD	Republic of Moldova	UA	Ukraine
ES	Spain	MG	Madagascar	UG	Uganda
FI	Finland	ML	Mali	US	United States of America
FR	France	MN	Mongolia	UZ	Uzbekistan
GA	Gabon	MR	Mauritania	VN	Viet Nam

- 1 -

**TITLE OF THE INVENTION****INHIBITORS OF FARNESYL-PROTEIN TRANSFERASE****BACKGROUND OF THE INVENTION**

5           The Ras protein is part of a signalling pathway that links cell surface growth factor receptors to nuclear signals initiating cellular proliferation. Biological and biochemical studies of Ras action indicate that Ras functions like a G-regulatory protein. In the inactive state, Ras is bound to GDP. Upon growth factor receptor activation Ras is induced  
10 to exchange GDP for GTP and undergoes a conformational change. The GTP-bound form of Ras propagates the growth stimulatory signal until the signal is terminated by the intrinsic GTPase activity of Ras, which returns the protein to its inactive GDP bound form (D.R. Lowy and D.M. Willumsen, *Ann. Rev. Biochem.* 62:851-891 (1993)). Mutated *ras* genes  
15 are found in many human cancers, including colorectal carcinoma, exocrine pancreatic carcinoma, and myeloid leukemias. The protein products of these genes are defective in their GTPase activity and constitutively transmit a growth stimulatory signal.

Ras must be localized to the plasma membrane for both  
20 normal and oncogenic functions. At least 3 post-translational modifications are involved with Ras membrane localization, and all 3 modifications occur at the C-terminus of Ras. The Ras C-terminus contains a sequence motif termed a "CAAX" or "Cys-Aaa<sup>1</sup>-Aaa<sup>2</sup>-Xaa" box (Cys is cysteine, Aaa is an aliphatic amino acid, the Xaa is any amino  
25 acid) (Willumsen *et al.*, *Nature* 310:583-586 (1984)). Depending on the specific sequence, this motif serves as a signal sequence for the enzymes farnesyl-protein transferase or geranylgeranyl-protein transferase, which catalyze the alkylation of the cysteine residue of the CAAX motif with a C<sub>15</sub> or C<sub>20</sub> isoprenoid, respectively. (S. Clarke., *Ann. Rev. Biochem.*  
30 61:355-386 (1992); W.R. Schafer and J. Rine, *Ann. Rev. Genetics* 30:209-237 (1992)). The Ras protein is one of several proteins that are known to undergo post-translational farnesylation. Other farnesylated proteins include the Ras-related GTP-binding proteins such as Rho, fungal mating factors, the nuclear lamins, and the gamma subunit of

- 2 -

transducin. James, et al., *J. Biol. Chem.* 269, 14182 (1994) have identified a peroxisome associated protein Pxf which is also farnesylated. James, et al., have also suggested that there are farnesylated proteins of unknown structure and function in addition to those listed above

5                   Inhibition of farnesyl-protein transferase has been shown to block the growth of Ras-transformed cells in soft agar and to modify other aspects of their transformed phenotype. It has also been demonstrated that certain inhibitors of farnesyl-protein transferase selectively block the processing of the Ras oncoprotein intracellularly  
10 (N.E. Kohl *et al.*, *Science*, 260:1934-1937 (1993) and G.L. James *et al.*, *Science*, 260:1937-1942 (1993). Recently, it has been shown that an inhibitor of farnesyl-protein transferase blocks the growth of *ras*-dependent tumors in nude mice (N.E. Kohl *et al.*, *Proc. Natl. Acad. Sci U.S.A.*, 91:9141-9145 (1994).

15                   Indirect inhibition of farnesyl-protein transferase *in vivo* has been demonstrated with lovastatin (Merck & Co., Rahway, NJ) and compactin (Hancock *et al.*, *ibid*; Casey *et al.*, *ibid*; Schafer *et al.*, *Science* 245:379 (1989)). These drugs inhibit HMG-CoA reductase, the rate limiting enzyme for the production of polyisoprenoids including farnesyl  
20 pyrophosphate. Farnesyl-protein transferase utilizes farnesyl pyrophosphate to covalently modify the Cys thiol group of the Ras CAAX box with a farnesyl group (Reiss *et al.*, *Cell*, 62:81-88 (1990); Schaber *et al.*, *J. Biol. Chem.*, 265:14701-14704 (1990); Schafer *et al.*, *Science*, 249:1133-1139 (1990); Manne *et al.*, *Proc. Natl. Acad. Sci USA*,  
25 87:7541-7545 (1990)). Inhibition of farnesyl pyrophosphate biosynthesis by inhibiting HMG-CoA reductase blocks Ras membrane localization in cultured cells. However, direct inhibition of farnesyl-protein transferase would be more specific and attended by fewer side effects than would occur with the required dose of a general inhibitor of isoprene  
30 biosynthesis.

                  Inhibitors of farnesyl-protein transferase (FPTase) have been described in two general classes. The first are analogs of farnesyl diphosphate (FPP), while the second class of inhibitors is related to the protein substrates (e.g., Ras) for the enzyme. In the peptide derived class



- 3 -

of inhibitors, a subclass of inhibitors has been described which generally comprises cysteine containing molecules that are related to the CAAX motif that is the signal for protein prenylation. (Schaber *et al.*, *ibid*; Reiss *et al.*, *ibid*; Reiss *et al.*, *PNAS*, 88:732-736 (1991)). Such inhibitors may  
5 inhibit protein prenylation while serving as alternate substrates for the farnesyl-protein transferase enzyme, or may be purely competitive inhibitors (U.S. Patent 5,141,851, University of Texas; N.E. Kohl *et al.*, *Science*, 260:1934-1937 (1993); Graham, *et al.*, *J. Med. Chem.*, 37, 725 (1994)). In general, deletion of the thiol from a peptidyl CAAX  
10 derivative has been shown to dramatically reduce the inhibitory potency of the compound. However, the thiol group potentially places limitations on the therapeutic application of FPTase inhibitors with respect to pharmacokinetics, pharmacodynamics and toxicity. Therefore, a functional replacement for the thiol is desirable. With the exception of a  
15 group of antibiotics known as the peptidocinnamins (Omura, *et al.*, *J. Antibiotics* 46:222 (1993)), non-thiol FPTase inhibitors that are competitive with the Ras substrate have not been described.

Another subclass of the peptide derived inhibitors which comprises peptidomimetic compounds wherein the central AA portion of  
20 the CAAX motif has been replaced by 3-aminobenzoic acid and 3-aminomethylbenzoic acid spacers has recently been described (M. Nigam *et al. J. Biol. Chem.*, 268:20695-20698 (1993), Y. Qian *et al. J. Biol. Chem.*, 269:12410-12413 (1994)). FPTase peptidomimetic inhibitors further lacking a C-terminus peptidyl moiety (wherein the X peptide has  
25 been replaced by a non-peptide moiety) have also been recently described (A. Vogt *et al. J. Biol. Chem.*, 270:660-664 (1995)). All of the compounds in this second subclass of peptide derived inhibitors retain the thiol moiety.

It is, therefore, an object of this invention to develop non-  
30 peptide compounds that do not have a thiol moiety, and that will inhibit farnesyl-protein transferase and thus, the post-translational farnesylation of proteins. It is a further object of this invention to develop chemotherapeutic compositions containing the compounds of this invention and methods for producing the compounds of this invention.





- 4 -

**SUMMARY OF THE INVENTION**

- The present invention includes substituted aminoalkylbenzamide and aminobenzamide analogs which inhibit the farnesyl-protein transferase, chemotherapeutic compositions containing the compounds of this invention, and methods for producing the compounds of this invention. Furthermore these analogs differ from those previously described as inhibitors of farnesyl-protein transferase in that they do not have a thiol moiety. The lack of the thiol offers unique advantages in terms of improved pharmacokinetic behavior in animals, prevention of thiol-dependent chemical reactions, such as rapid autoxidation and disulfide formation with endogenous thiols, and reduced systemic toxicity.

- 5 -

The compounds of this invention are illustrated by the formulae:




5

### DETAILED DESCRIPTION OF THE INVENTION

The compounds of this invention are useful in the inhibition of farnesyl-protein transferase and the farnesylation of certain proteins.

- 6 -

In a first embodiment of this invention, the farnesyl-protein transferase inhibitors are illustrated by the formula I:



5 wherein:

R1a and R1b are independently selected from:

- a) hydrogen,
- b) aryl, heterocycle, cycloalkyl, alkenyl, alkynyl, R10O-,  
 10 R11S(O)m-, R10C(O)NR10-, CN, NO2,  
 (R10)2N-C(NR10)-, R10C(O)-, R10OC(O)-, N3, -N(R10)2,  
 or R11OC(O)NR10-,
- c) C1-C6 alkyl unsubstituted or substituted by aryl,  
 heterocyclic, cycloalkyl, alkenyl, alkynyl, R10O-,  
 R11S(O)m-, R10C(O)NR10-, CN, (R10)2N-C(NR10)-,  
 15 R10C(O)-, R10OC(O)-, N3, -N(R10)2, or R11OC(O)-  
 NR10-;

R2a and R2b are independently selected from:

- a) hydrogen,
- b) C1-C6 alkyl unsubstituted or substituted by alkenyl, R10O-,  
 20 R11S(O)m-, R10C(O)NR10-, CN, N3, (R10)2N-C(NR10)-,  
 R10C(O)-, R10OC(O)-, -N(R10)2, or R11OC(O)NR10-,
- c) aryl, heterocycle, cycloalkyl, alkenyl, R10O-,  
 R11S(O)m-, R10C(O)NR10-, CN, NO2,  
 25 (R10)2N-C(NR10)-, R10C(O)-, R10OC(O)-, N3,  
 -N(R10)2, or R11OC(O)NR10-, and
- d) C1-C6 alkyl substituted with an unsubstituted or  
 substituted group selected from aryl, heterocyclic and  
 C3-C10 cycloalkyl;

30

- 7 -

R3a and R3b are independently selected from:

- a) a side chain of a naturally occurring amino acid,
- b) an oxidized form of a side chain of a naturally occurring amino acid which is:
  - i) methionine sulfoxide, or
  - ii) methionine sulfone, and
- c) substituted or unsubstituted C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, aryl or heterocyclic group,
 

wherein the substituent is selected from F, Cl, Br, N(R<sup>10</sup>)<sub>2</sub>, NO<sub>2</sub>, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>-, -N(R<sup>10</sup>)<sub>2</sub>, R<sup>11</sup>OC(O)NR<sup>10</sup>- and C<sub>1</sub>-C<sub>20</sub> alkyl, and
- d) C<sub>1</sub>-C<sub>6</sub> alkyl substituted with an unsubstituted or substituted group selected from aryl, heterocycle and C<sub>3</sub>-C<sub>10</sub> cycloalkyl; or

R3a and R3b are combined to form -(CH<sub>2</sub>)<sub>s</sub> - wherein one of the carbon atoms is optionally replaced by a moiety selected from: O, S(O)<sub>m</sub>, -NC(O)-, and -N(COR<sup>10</sup>)- ;

R4 and R5 are independently selected from:

- a) hydrogen, and
- b)
 
$$\begin{array}{c}
 (R^7)_r \\
 | \\
 V - A^1(CR^{1a}_2)_n A^2(CR^{1a}_2)_n - \left( \begin{array}{c} R^8 \\ | \\ W \end{array} \right)_u - (CR^{1b}_2)_p - \begin{array}{c} Z \\ || \\ \text{---} \end{array}
 \end{array}
 ;$$

R<sup>7</sup> is independently selected from:

- a) hydrogen,
- b) aryl, heterocycle, cycloalkyl, alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, R<sup>10</sup><sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and

- 8 -

- 5 c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl, heterocycle, cycloalkyl, alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NH-, CN, H<sub>2</sub>N-C(NH)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>10</sup>OC(O)NH-;

R<sup>8</sup> is selected from:

- 10 a) hydrogen,  
b) alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C-(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and  
15 c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by perfluoroalkyl, F, Cl, Br, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-;

R<sup>10</sup> is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl and aryl;

20

R<sup>11</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

25 A<sup>1</sup> and A<sup>2</sup> are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR<sup>10</sup>-, -NR<sup>10</sup>C(O)-, O, -N(R<sup>10</sup>)-, -S(O)<sub>2</sub>N(R<sup>10</sup>)-, -N(R<sup>10</sup>)S(O)<sub>2</sub>-, or S(O)<sub>m</sub>;

25

V is selected from:

- 30 a) hydrogen,  
b) heterocycle,  
c) aryl,  
d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and  
e) C<sub>2</sub>-C<sub>20</sub> alkenyl,

- 9 -

provided that V is not hydrogen if  $A^1$  is  $S(O)_m$  and V is not hydrogen if  $A^1$  is a bond, n is 0 and  $A^2$  is  $S(O)_m$ ;

W is a heterocycle;

5

Z is independently  $H_2$  or O;

m is 0, 1 or 2;

n is 0, 1, 2, 3 or 4;

10 p is 0, 1, 2, 3 or 4;

r is 0 to 5, provided that r is 0 when V is hydrogen;


s is 4 or 5; and

u is 0 or 1;

or the pharmaceutically acceptable salts thereof.

15

In a second embodiment of this invention the prodrugs of compounds of formula I are illustrated by the formula II:



20 wherein:

$R^{1a}$  and  $R^{1b}$  are independently selected from:

a) hydrogen,

b) aryl, heterocycle, cycloalkyl, alkenyl, alkynyl,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $NO_2$ ,

25  $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ ,

c)  $C_1-C_6$  alkyl unsubstituted or substituted by aryl, heterocyclic, cycloalkyl, alkenyl, alkynyl,  $R^{10}O-$ ,

- 10 -

$R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $(R^{10})_2N-C(NR^{10})-$ ,  
 $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)-$   
 $NR^{10}-$ ;

5 R2a and R2b are independently selected from:

- a) hydrogen,
- b) C1-C6 alkyl unsubstituted or substituted by alkenyl,  $R^{10}O-$ ,  
 $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $N_3$ ,  $(R^{10})_2N-C(NR^{10})-$ ,  
 $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ ,
- 10 c) aryl, heterocycle, cycloalkyl, alkenyl,  $R^{10}O-$ ,  
 $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $NO_2$ ,  $(R^{10})_2N-$   
 $C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ ,  
or  $R^{11}OC(O)NR^{10}-$ , and
- d) C1-C6 alkyl substituted with an unsubstituted or  
15 substituted group selected from aryl, heterocyclic and  
C3- C10 cycloalkyl;

R3a and R3b are independently selected from:

- 20 a) a side chain of a naturally occurring amino acid,
- b) an oxidized form of a side chain of a naturally occurring  
amino acid which is:
  - i) methionine sulfoxide, or
  - ii) methionine sulfone, and
- 25 c) substituted or unsubstituted C1-C20 alkyl, C2-C20 alkenyl,  
C3-C10 cycloalkyl, aryl or heterocyclic group,  
wherein the substituent is selected from F, Cl, Br,  
 $N(R^{10})_2$ ,  $NO_2$ ,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ ,  
CN,  $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  
30  $-N(R^{10})_2$ ,  $R^{11}OC(O)NR^{10}-$  and C1-C20 alkyl, and
- d) C1-C6 alkyl substituted with an unsubstituted or  
substituted group selected from aryl, heterocycle and C3-  
C10 cycloalkyl; or




- 11 -

R3a and R3b are combined to form  $-(CH_2)_s-$  wherein one of the carbon atoms is optionally replaced by a moiety selected from: O, S(O)<sub>m</sub>, -NC(O)-, and -N(COR<sup>10</sup>)-;

5 R<sup>4</sup> and R<sup>5</sup> are independently selected from:

- a) hydrogen, and
- b)




10 R<sup>6</sup> is

- a) substituted or unsubstituted C<sub>1</sub>-C<sub>8</sub> alkyl or substituted or unsubstituted C<sub>5</sub>-C<sub>8</sub> cycloalkyl, wherein the substituent on the alkyl is selected from:
  - 1) aryl,
  - 2) heterocycle,
  - 3) -N(R<sup>11</sup>)<sub>2</sub>,
  - 4) -OR<sup>10</sup>, or

15

b)



20

R<sup>7</sup> is independently selected from:

- a) hydrogen,
- b) aryl, heterocycle, cycloalkyl, alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, R<sup>10</sup><sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and
- c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl, heterocycle, cycloalkyl, alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NH-, CN, H<sub>2</sub>N-

25

- 12 -

C(NH)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>10</sup>OC(O)NH-;

R<sup>8</sup> is selected from:

- 5           a) hydrogen,
- b) alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C-(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and
- 10          c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by perfluoroalkyl, F, Cl, Br, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, (R<sup>10</sup>)<sub>2</sub>N-C-(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-;

- 15       R<sup>10</sup> is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl and aryl;

R<sup>11</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

- 20       R<sup>12</sup> is independently selected from hydrogen and C<sub>1</sub>-C<sub>6</sub> alkyl;

R<sup>13</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl;

- 25       A<sup>1</sup> and A<sup>2</sup> are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR<sup>10</sup>-, -NR<sup>10</sup>C(O)-, O, -N(R<sup>10</sup>)-, -S(O)<sub>2</sub>N(R<sup>10</sup>)-, -N(R<sup>10</sup>)S(O)<sub>2</sub>-, or S(O)<sub>m</sub>;

V is selected from:

- 30           a) hydrogen,
- b) heterocycle,
- c) aryl,
- d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
- e) C<sub>2</sub>-C<sub>20</sub> alkenyl,

- 13 -

provided that V is not hydrogen if  $A^1$  is  $S(O)_m$  and V is not hydrogen if  $A^1$  is a bond, n is 0 and  $A^2$  is  $S(O)_m$ ;

W is a heterocycle;

5

Z is independently  $H_2$  or O;

m is 0, 1 or 2;

n is 0, 1, 2, 3 or 4;

10 p is 0, 1, 2, 3 or 4;


r is 0 to 5, provided that r is 0 when V is hydrogen;

s is 4 or 5; and

u is 0 or 1;

15 or the pharmaceutically acceptable salts thereof.

In a third embodiment of this invention, the inhibitors of farnesyl transferase are illustrated by the formula III:



20 wherein:

$R^{1a}$  and  $R^{1b}$  are independently selected from:

a) hydrogen,

b) aryl, heterocycle, cycloalkyl, alkenyl, alkynyl,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $NO_2$ ,

25  $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ ,

- 14 -


- 5 c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl, heterocyclic, cycloalkyl, alkenyl, alkynyl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)-NR<sup>10</sup>-;

R<sup>2a</sup> and R<sup>2b</sup> are independently selected from:

- 10 a) hydrogen,  
 b) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by alkenyl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, N<sub>3</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-,  
 15 c) aryl, heterocycle, cycloalkyl, alkenyl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and  
 d) C<sub>1</sub>-C<sub>6</sub> alkyl substituted with an unsubstituted or substituted group selected from aryl, heterocyclic and C<sub>3</sub>-C<sub>10</sub> cycloalkyl;

20 R<sup>4</sup> and R<sup>5</sup> are independently selected from:

- a) hydrogen, and  
 b)



25 R<sup>7</sup> is independently selected from:

- a) hydrogen,  
 b) aryl, heterocycle, cycloalkyl, alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, R<sup>10</sup><sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and  
 30

- 15 -

- 5 c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl, heterocycle, cycloalkyl, alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NH-, CN, H<sub>2</sub>N-C(NH)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>10</sup>OC(O)NH-;

R<sup>8</sup> is selected from:

- 10 a) hydrogen,  
b) alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C-(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and  
15 c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by perfluoroalkyl, F, Cl, Br, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-;

20 R<sup>10</sup> is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl and aryl;

R<sup>11</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

25 A<sup>1</sup> and A<sup>2</sup> are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR<sup>10</sup>-, -NR<sup>10</sup>C(O)-, O-, -N(R<sup>10</sup>)-, -S(O)<sub>2</sub>N(R<sup>10</sup>)-, -N(R<sup>10</sup>)S(O)<sub>2</sub>-, or S(O)<sub>m</sub>;

V is selected from:

- 30 a) hydrogen,  
b) heterocycle,  
c) aryl,  
d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and  
e) C<sub>2</sub>-C<sub>20</sub> alkenyl,

- 16 -

provided that V is not hydrogen if  $A^1$  is  $S(O)_m$  and V is not hydrogen if  $A^1$  is a bond, n is 0 and  $A^2$  is  $S(O)_m$ ;

W is a heterocycle;

5


Z is independently  $H_2$  or O;

- m is 0, 1 or 2;  
 n is 0, 1, 2, 3 or 4;  
 10 p is 0, 1, 2, 3 or 4;  
 q is 0, 1 or 2;  
 r is 0 to 5, provided that r is 0 when V is hydrogen;  
 s is 4 or 5; and  
 u is 0 or 1;

15

or the pharmaceutically acceptable salts thereof.

In a fourth embodiment of this invention the prodrugs of compounds of formula III are illustrated by the formula IV:



20

wherein:

$R^{1a}$  and  $R^{1b}$  are independently selected from:

- 25 a) hydrogen,  
 b) aryl, heterocycle, cycloalkyl, alkenyl, alkynyl,  $R^{10}O-$ ,  
 $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $NO_2$ ,  
 $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ ,  
 or  $R^{11}OC(O)NR^{10}-$ ,

- 17 -


- 5 c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl, heterocyclic, cycloalkyl, alkenyl, alkynyl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-;

R<sup>2a</sup> and R<sup>2b</sup> are independently selected from:

- 10 a) hydrogen,  
 b) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by alkenyl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, N<sub>3</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-,  
 15 c) aryl, heterocycle, cycloalkyl, alkenyl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and  
 d) C<sub>1</sub>-C<sub>6</sub> alkyl substituted with an unsubstituted or substituted group selected from aryl, heterocyclic and C<sub>3</sub>-C<sub>10</sub> cycloalkyl;

20 R<sup>4</sup> and R<sup>5</sup> are independently selected from:

- a) hydrogen, and  
 b)



25 R<sup>7</sup> is independently selected from:

- a) hydrogen,  
 b) aryl, heterocycle, cycloalkyl, alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, R<sup>10</sup><sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, 30 R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and  
 c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl, heterocycle, cycloalkyl, alkenyl, alkynyl, perfluoroalkyl, F,

- 18 -

Cl, Br,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NH-$ , CN,  $H_2N-$ ,  $C(NH)-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{10}OC(O)NH-$ ;

5 R<sup>8</sup> is selected from:

- a) hydrogen,
- b) alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN, NO<sub>2</sub>,  $(R^{10})_2N-C-(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ , and
- 10 c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by perfluoroalkyl, F, Cl, Br,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $(R^{10})_2N-C-(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ ;

15

R<sup>10</sup> is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl and aryl;

R<sup>11</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

20

A<sup>1</sup> and A<sup>2</sup> are independently selected from: a bond,  $-CH=CH-$ ,  $-C\equiv C-$ ,  $-C(O)-$ ,  $-C(O)NR^{10}-$ ,  $-NR^{10}C(O)-$ , O,  $-N(R^{10})-$ ,  $-S(O)_2N(R^{10})-$ ,  $-N(R^{10})S(O)_2-$ , or  $S(O)_m$ ;

25 V is selected from:

- a) hydrogen,
- b) heterocycle,
- c) aryl,
- d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are replaced
- 30 with a heteroatom selected from O, S, and N, and
- e) C<sub>2</sub>-C<sub>20</sub> alkenyl,

provided that V is not hydrogen if A<sup>1</sup> is  $S(O)_m$  and V is not hydrogen if A<sup>1</sup> is a bond, n is 0 and A<sup>2</sup> is  $S(O)_m$ ;



- 19 -


W is a heterocycle;

Z is independently H<sub>2</sub> or O;

- 5    m is            0, 1 or 2;  
      n is            0, 1, 2, 3 or 4;  
      p is            0, 1, 2, 3 or 4;  
      q is            0, 1 or 2;  
      r is            0 to 5, provided that r is 0 when V is hydrogen;  
 10    s is            4 or 5; and  
      u is            0 or 1;

or the pharmaceutically acceptable salts thereof.

- 15                    In a more preferred embodiment of this invention, the Ras farnesyl transferase inhibitors are illustrated by the Formula Ia:



- 20    wherein:

R<sup>1a</sup> is independently selected from: hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl;

R<sup>1b</sup> is independently selected from:

- 25            a)    hydrogen,  
              b)    aryl, heterocycle, cycloalkyl, R<sup>10</sup>O-, -N(R<sup>10</sup>)<sub>2</sub> or alkenyl,  
              c)    C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl,  
                      heterocycle, cycloalkyl, alkenyl, R<sup>10</sup>O-, or -N(R<sup>10</sup>)<sub>2</sub>;

R<sup>2a</sup> is selected from:

- 30            a)    hydrogen,

- 20 -

- 5           b) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by alkenyl, R<sup>10</sup>O-,  
R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, N<sub>3</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-,  
R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-,  
c) aryl, heterocycle, cycloalkyl, alkenyl, R<sup>10</sup>O-,  
R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>,  
10           (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>,  
-N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and  
d) C<sub>1</sub>-C<sub>6</sub> alkyl substituted with an unsubstituted or  
substituted group selected from aryl, heterocyclic and  
10       C<sub>3</sub>- C<sub>10</sub> cycloalkyl;

R<sub>2b</sub> is hydrogen;


R<sub>3a</sub> and R<sub>3b</sub> are independently selected from:

- 15           a) a side chain of a naturally occurring amino acid,  
b) an oxidized form of a side chain of a naturally occurring  
amino acid which is:  
i) methionine sulfoxide, or  
ii) methionine sulfone,  
20           c) substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>2</sub>-C<sub>10</sub> alkenyl,  
C<sub>3</sub>-C<sub>10</sub> cycloalkyl, aryl or heterocyclic group,  
wherein the substituent is selected from F, Cl, Br,  
NO<sub>2</sub>, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN,  
(R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>,  
25           -N(R<sup>10</sup>)<sub>2</sub>, R<sup>11</sup>OC(O)NR<sup>10</sup>- and C<sub>1</sub>-C<sub>20</sub> alkyl, and  
d) C<sub>1</sub>-C<sub>6</sub> alkyl substituted with an unsubstituted or  
substituted group selected from aryl, heterocycle and C<sub>3</sub>-  
C<sub>10</sub> cycloalkyl;

30       R<sub>4</sub> and R<sub>5</sub> are independently selected from:

- a) hydrogen, and  
b)

- 21 -



$R^7$  is independently selected from:

- a) hydrogen,
- 5 b) C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and
- 10 c) C<sub>1</sub>-C<sub>6</sub> alkyl substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>10</sup>O-, R<sup>10</sup>C(O)NR<sup>10</sup>-, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-;

$R^8$  is selected from:

- a) hydrogen,
- 15 b) C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and
- 20 c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-;

25  $R^{10}$  is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl and aryl;

$R^{11}$  is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

30  $A^1$  and  $A^2$  are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR<sup>10</sup>-, O, -N(R<sup>10</sup>)-, or S(O)<sub>m</sub>;

V is selected from:

- 22 -

- a) hydrogen,  
 b) heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, isoquinolinyl, and thienyl,  
 5 c) aryl,  
 d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and  
 e) C<sub>2</sub>-C<sub>20</sub> alkenyl, and  
 provided that V is not hydrogen if A<sup>1</sup> is S(O)<sub>m</sub> and V is not hydrogen if  
 10 A<sup>1</sup> is a bond, n is 0 and A<sup>2</sup> is S(O)<sub>m</sub>;

W is a heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, or isoquinolinyl;

15


Z is independently H<sub>2</sub> or O;

- m is 0, 1 or 2;  
 n is 0, 1, 2, 3 or 4;  
 20 p is 0, 1, 2, 3 or 4;  
 r is 0 to 5, provided that r is 0 when V is hydrogen; and  
 u is 0 or 1;

or the pharmaceutically acceptable salts thereof.

25

In a second more preferred embodiment of this invention, the prodrugs of the preferred compounds of Formula I are illustrated by the Formula IIa:



- 23 -

R1a is independently selected from: hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl;

R1b is independently selected from:

- 5       a) hydrogen,
- b) aryl, heterocycle, cycloalkyl, R<sup>10</sup>O-, -N(R<sup>10</sup>)<sub>2</sub> or alkenyl,
- c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl,  
             heterocycle, cycloalkyl, alkenyl, R<sup>10</sup>O-, or -N(R<sup>10</sup>)<sub>2</sub>;
- 10      R2a is selected from:
  - a) hydrogen,
  - b) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by alkenyl, R<sup>10</sup>O-,  
R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, N<sub>3</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-,  
R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-,
  - 15       c) aryl, heterocycle, cycloalkyl, alkenyl, R<sup>10</sup>O-,  
R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-  
C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>,  
or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and
  - d) C<sub>1</sub>-C<sub>6</sub> alkyl substituted with an unsubstituted or
  - 20       substituted group selected from aryl, heterocyclic and  
C<sub>3</sub>- C<sub>10</sub> cycloalkyl;

R2b is hydrogen;

25      R3a and R3b are independently selected from:

- a) a side chain of a naturally occurring amino acid,
- b) an oxidized form of a side chain of a naturally occurring  
amino acid which is:
  - i) methionine sulfoxide, or
  - 30       ii) methionine sulfone,
- c) substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>2</sub>-C<sub>10</sub> alkenyl,  
C<sub>3</sub>-C<sub>10</sub> cycloalkyl, aryl or heterocyclic group,  
wherein the substituent is selected from F, Cl, Br,  
NO<sub>2</sub>, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN,

- 24 -


$(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  
 $-N(R^{10})_2$ ,  $R^{11}OC(O)NR^{10}-$  and  $C_1-C_{20}$  alkyl, and

- d)  $C_1-C_6$  alkyl substituted with an unsubstituted or substituted group selected from aryl, heterocycle and  $C_3-C_{10}$  cycloalkyl;

$R^4$  and  $R^5$  are independently selected from:

a) hydrogen, and

b)




$R^6$  is

- a) substituted or unsubstituted  $C_1-C_8$  alkyl or substituted or unsubstituted  $C_5-C_8$  cycloalkyl, wherein the substituent on the alkyl is selected from:

- 1) aryl,
- 2) heterocycle,
- 3)  $-N(R^{11})_2$ ,
- 4)  $-OR^{10}$ , or

b)



$R^7$  is independently selected from:

- a) hydrogen,
- b)  $C_1-C_6$  alkyl,  $C_2-C_6$  alkenyl,  $C_2-C_6$  alkynyl,  $C_1-C_6$  perfluoroalkyl, F, Cl,  $R^{10}O-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $NO_2$ ,  $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ , and
- c)  $C_1-C_6$  alkyl substituted by  $C_1-C_6$  perfluoroalkyl,  $R^{10}O-$ ,  $R^{10}C(O)NR^{10}-$ ,  $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ ;

- 25 -

R<sup>8</sup> is selected from:

- 5           a) hydrogen,  
          b) C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and  
          c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-;
- 10

R<sup>10</sup> is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl and aryl;

15

R<sup>11</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

R<sup>12</sup> is independently selected from hydrogen and C<sub>1</sub>-C<sub>6</sub> alkyl;

20 R<sup>13</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl;

A<sup>1</sup> and A<sup>2</sup> are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR<sup>10</sup>-, O, -N(R<sup>10</sup>)-, or S(O)<sub>m</sub>;

25 V is selected from:

- a) hydrogen,  
          b) heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, isoquinolinyl, and thienyl,  
30           c) aryl,  
          d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and  
          e) C<sub>2</sub>-C<sub>20</sub> alkenyl, and

- 26 -

provided that V is not hydrogen if  $A^1$  is  $S(O)_m$  and V is not hydrogen if  $A^1$  is a bond, n is 0 and  $A^2$  is  $S(O)_m$ ;


W is a heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl,  
5 thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, or  
isoquinolinyl;

Z is independently  $H_2$  or O;

10 m is 0, 1 or 2;  
n is 0, 1, 2, 3 or 4;  
p is 0, 1, 2, 3 or 4;  
r is 0 to 5, provided that r is 0 when V is hydrogen; and  
u is 0 or 1;

15 or the pharmaceutically acceptable salts thereof.

In a third more preferred embodiment of this invention, the inhibitors of farnesyl transferase are illustrated by the formula IIIa:



IIIa

wherein:

$R^{1a}$  is independently selected from: hydrogen or  $C_1$ - $C_6$  alkyl;

$R^{1b}$  is independently selected from:

25 a) hydrogen,  
b) aryl, heterocycle, cycloalkyl,  $R^{10}O$ -,  $-N(R^{10})_2$  or alkenyl,



- 27 -

- c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl, heterocycle, cycloalkyl, alkenyl, R<sup>10</sup>O-, or -N(R<sup>10</sup>)<sub>2</sub>;


R<sub>2a</sub> is selected from:

- 5 a) hydrogen,  
 b) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by alkenyl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, N<sub>3</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-,  
 10 c) aryl, heterocycle, cycloalkyl, alkenyl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and  
 d) C<sub>1</sub>-C<sub>6</sub> alkyl substituted with an unsubstituted or substituted group selected from aryl, heterocyclic and  
 15 C<sub>3</sub>- C<sub>10</sub> cycloalkyl;

R<sub>2b</sub> is hydrogen;

R<sub>4</sub> and R<sub>5</sub> are independently selected from:

- 20 a) hydrogen, and  
 b)



R<sub>7</sub> is independently selected from:

- 25 a) hydrogen,  
 b) C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and  
 30 c) C<sub>1</sub>-C<sub>6</sub> alkyl substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>10</sup>O-, R<sup>10</sup>C(O)NR<sup>10</sup>-, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-;

- 28 -

R<sup>8</sup> is selected from:

- 5           a) hydrogen,  
          b) C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and  
10           c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-;

R<sup>10</sup> is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl and aryl;

15

R<sup>11</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

A<sup>1</sup> and A<sup>2</sup> are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR<sup>10</sup>-, O, -N(R<sup>10</sup>)-, or S(O)<sub>m</sub>;

20

V is selected from:

- a) hydrogen,  
          b) heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, isoquinolinyl, and thienyl,  
25           c) aryl,  
          d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and  
          e) C<sub>2</sub>-C<sub>20</sub> alkenyl, and  
30   provided that V is not hydrogen if A<sup>1</sup> is S(O)<sub>m</sub> and V is not hydrogen if A<sup>1</sup> is a bond, n is 0 and A<sup>2</sup> is S(O)<sub>m</sub>;

- 29 -

W is a heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, or isoquinolinyl;

5 Z is independently H<sub>2</sub> or O;

m is 0, 1 or 2;

n is 0, 1, 2, 3 or 4;

p is 0, 1, 2, 3 or 4;

10 q is 0, 1 or 2;


r is 0 to 5, provided that r is 0 when V is hydrogen; and

u is 0 or 1;

or the pharmaceutically acceptable salts thereof.

15

In a fourth more preferred embodiment of this invention, the prodrugs of the preferred compounds of Formula III are illustrated by the Formula IVa:



20

wherein:

R<sup>1a</sup> is independently selected from: hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl;

R<sup>1b</sup> is independently selected from:

25

- a) hydrogen,
- b) aryl, heterocycle, cycloalkyl, R<sup>10</sup>O-, -N(R<sup>10</sup>)<sub>2</sub> or alkenyl,
- c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl, heterocycle, cycloalkyl, alkenyl, R<sup>10</sup>O-, or -N(R<sup>10</sup>)<sub>2</sub>;

- 30 -


R2a is selected from:

- 5 a) hydrogen,  
 b) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by alkenyl, R<sup>10</sup>O-,  
 R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, N<sub>3</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-,  
 R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-,  
 c) aryl, heterocycle, cycloalkyl, alkenyl, R<sup>10</sup>O-,  
 R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>,  
 (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-,  
 10 N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and  
 d) C<sub>1</sub>-C<sub>6</sub> alkyl substituted with an unsubstituted or  
 substituted group selected from aryl, heterocyclic and  
 C<sub>3</sub>- C<sub>10</sub> cycloalkyl;

15 R2b is hydrogen;

R4 and R5 are independently selected from:

- a) hydrogen, and  
 b)



R7 is independently selected from:

- 25 a) hydrogen,  
 b) C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub>  
 perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>,  
 (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or  
 R<sup>11</sup>OC(O)NR<sup>10</sup>-, and  
 c) C<sub>1</sub>-C<sub>6</sub> alkyl substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>10</sup>O-,  
 R<sup>10</sup>C(O)NR<sup>10</sup>-, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-,  
 30 R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-;

R8 is selected from:

- 31 -

- 5
- a) hydrogen,
  - b) C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and
  - c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-;

10 R<sup>10</sup> is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl and aryl;

15 R<sup>11</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

15 A<sup>1</sup> and A<sup>2</sup> are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR<sup>10</sup>-, O, -N(R<sup>10</sup>)-, or S(O)<sub>m</sub>;

V is selected from:

- 20
- a) hydrogen,
  - b) heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, isoquinolinyl, and thienyl,
  - c) aryl,
  - 25 d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
  - e) C<sub>2</sub>-C<sub>20</sub> alkenyl, and

provided that V is not hydrogen if A<sup>1</sup> is S(O)<sub>m</sub> and V is not hydrogen if A<sup>1</sup> is a bond, n is 0 and A<sup>2</sup> is S(O)<sub>m</sub>;

30

W is a heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, or isoquinolinyl;

- 32 -

Z is independently H<sub>2</sub> or O;

- m is 0, 1 or 2;  
n is 0, 1, 2, 3 or 4;  
5 p is 0, 1, 2, 3 or 4;  
q is 0, 1 or 2;  
r is 0 to 5, provided that r is 0 when V is hydrogen; and  
u is 0 or 1;

- 10 or the pharmaceutically acceptable salts thereof.

The preferred compounds of this invention are as follows:

- 15 N-(1(S)-carboxy-3-methylthiopropyl)-3-(4-imidazolylmethyl)-  
aminomethylbenzamide

N-(1(S)-carbomethoxy-3-methylthiopropyl)-3-(4-imidazolylmethyl)-  
aminomethylbenzamide

- 20 N-(1(S)-carboxy-3-methylthiopropyl)-3-[N,N-*bis*-(4-imidazole-  
methyl)aminomethyl]benzamide

- N-(1(S)-carbomethoxy-3-methylthiopropyl)-3-[N,N-*bis*-(4-imidazole-  
methyl)aminomethyl]benzamide

- 25 N-(1(S)-carboxy-3-methylthiopropyl) 3-[(4-imidazolylmethyl) -N-  
methylaminomethyl]benzamide

- 30 N-(1(S)-carbomethoxy-3-methylthiopropyl) 3-[(4-imidazolylmethyl) -N-  
methylaminomethyl]benzamide

N-(1(S)-carboxy-3-methylthiopropyl) 4-[(4-imidazolylmethyl)-  
amino]benzamide

- 33 -

N-(1(S)-carbomethoxy-3-methylthiopropyl) 4-[(4-imidazolylmethyl)-  
amino]benzamide

5 N-(1(S)-carboxy-3-methylthiopropyl)-3-[(4-imidazolylmethyl) amino]  
benzamide

N-(1(S)-carbomethoxy-3-methylthiopropyl)-3-[(4-imidazolylmethyl)  
amino] benzamide

10 N-(1(S)-carbomethoxy-3-methylthiopropyl)-3-[(4-imidazolylpropyl)  
amino]benzamide

N-(1(S)-Carboxy-3-methylthiopropyl)-3-[(4-imidazolylpropyl) amino]  
benzamide

15 N-(1(S)-Carbomethoxy-3-methylthiopropyl)-3-[N-(4-  
imidazolylmethyl)-N-(4-nitrobenzyl)aminomethyl]benzamide

20 N-(1(S)-Carboxy-3-methylthiopropyl)-3-[N-(4-imidazolylmethyl)-N-(4-  
nitrobenzyl)aminomethyl]benzamide

N-(1(S)-carboxy-3-methylthiopropyl)-3-[N,N-bis-(4-  
nitrophenylmethyl)]aminomethylbenzamide


25 N-(1(S)-carbomethoxy-3-methylthiopropyl)-3-[N,N-bis-(4-  
nitrophenylmethyl)aminomethyl]benzamide

or the pharmaceutically acceptable salts thereof.


30 Specific examples of the compounds of the invention are:

N-(1(S)-carboxy-3-methylthiopropyl)-3-[N,N-bis-(4-  
nitrophenylmethyl)aminomethyl]benzamide

- 34 -




N-(1(S)-carboxy-3-methylthiopropyl)-3-[N,N-bis-(4-nitrophenylmethyl)aminomethyl]benzamide



5


N-(1(S)-carboxy-3-methylthiopropyl)-3-[N,N-bis(4-imidazole-methyl)aminomethyl] benzamide






- 35 -


N-(1(S)-carboxomethoxy-3-methylthiopropyl)-3-[N,N-bis(4-imidazole-methyl)aminomethyl] benzamide



5 N-(1(S)-Carboxy-3-methylthiopropyl)-3-[N-(4-imidazolylmethyl)-N-(4-nitrobenzyl)aminomethyl]benzamide



N-(1(S)-Carbomethoxy-3-methylthiopropyl)-3-[N-(4-imidazolylmethyl)-N-(4-nitrobenzyl)aminomethyl]benzamide



- 36 -

or the pharmaceutically acceptable salts thereof.

In the present invention, the amino acids which are disclosed are identified both by conventional 3 letter and single letter abbreviations  
5 as indicated below:

---

	Alanine	Ala	A
	Arginine	Arg	R
10	Asparagine	Asn	N
	Aspartic acid	Asp	D
	Asparagine or		
	Aspartic acid	Asx	B
	Cysteine	Cys	C
15	Glutamine	Gln	Q
	Glutamic acid	Glu	E
	Glutamine or		
	Glutamic acid	Glx	Z
	Glycine	Gly	G
20	Histidine	His	H
	Isoleucine	Ile	I
	Leucine	Leu	L
	Lysine	Lys	K
	Methionine	Met	M
25	Phenylalanine	Phe	F
	Proline	Pro	P
	Serine	Ser	S
	Threonine	Thr	T
	Tryptophan	Trp	W
30	Tyrosine	Tyr	Y
	Valine	Val	V

---

- 37 -

The compounds of the present invention may have asymmetric centers and occur as racemates, racemic mixtures, and as individual diastereomers, with all possible isomers, including optical isomers, being included in the present invention.

5           As used herein, "alkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms.

          As used herein, "cycloalkyl" is intended to include non-aromatic cyclic hydrocarbon groups having the specified number of  
10 carbon atoms. Examples of cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and the like.

          "Alkenyl" groups include those groups having the specified number of carbon atoms and having one or several double bonds. Examples of alkenyl groups include vinyl, allyl, isopropenyl, pentenyl,  
15 hexenyl, heptenyl, cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, 1-propenyl, 2-butenyl, 2-methyl-2-butenyl, isoprenyl, farnesyl, geranyl, geranylgeranyl and the like.

          As used herein, "aryl" is intended to include any stable monocyclic, bicyclic or tricyclic carbon ring(s) of up to 7 members in  
20 each ring, wherein at least one ring is aromatic. Examples of aryl groups include phenyl, naphthyl, anthracenyl, biphenyl, tetrahydronaphthyl, indanyl, phenanthrenyl and the like.

          The term heterocycle or heterocyclic, as used herein, represents a stable 5- to 7-membered monocyclic or stable 8- to 11-  
25 membered bicyclic or stable 11-15 membered tricyclic heterocycle ring which is either saturated or unsaturated, and which consists of carbon atoms and from one to four heteroatoms selected from the group consisting of N, O, and S, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The  
30 heterocyclic ring may be attached at any heteroatom or carbon atom which results in the creation of a stable structure. Examples of such heterocyclic elements include, but are not limited to, azepinyl, benzimidazolyl, benzisoxazolyl, benzofurazanyl, benzopyranyl, benzothiopyranyl, benzofuryl, benzothiazolyl, benzothienyl,

- 38 -

benzoxazolyl, chromanyl, cinnoliny, dihydrobenzofuryl, dihydro-  
 benzothienyl, dihydrobenzothiopyranyl, dihydrobenzothio-pyranyl  
 sulfone, furyl, imidazolidinyl, imidazoliny, imidazolyl, indoliny,  
 indolyl, isochromanyl, isoindoliny, isoquinoliny, isothiazolidinyl,  
 5 isothiazolyl, isothiazolidinyl, morpholiny, naphthyridiny, oxadiazolyl,  
 2-oxoazepiny, 2-oxopiperaziny, 2-oxopiperidiny, 2-oxopyrrolidiny,  
 piperidyl, piperaziny, pyridyl, pyridyl N-oxide, pyridonyl, pyraziny,  
 pyrazolidinyl, pyrazolyl, pyrimidinyl, pyrrolidiny, pyrrolyl, quinazoliny,  
 quinoliny, quinoliny N-oxide, quinoxaliny, tetrahydrofuryl,  
 10 tetrahydroisoquinoliny, tetrahydro-quinoliny, thiamorpholiny,  
 thiamorpholiny sulfoxide, thiazolyl, thiazoliny, thienofuryl,  
 thienothienyl, and thienyl.


As used herein, the terms "substituted aryl", "substituted  
 heterocycle" and "substituted cycloalkyl" are intended to include the  
 15 cyclic group which is substituted with 1 or 2 substituents selected from  
 the group which includes but is not limited to F, Cl, Br, CF<sub>3</sub>, NH<sub>2</sub>,  
 N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>, NO<sub>2</sub>, CN, (C<sub>1</sub>-C<sub>6</sub> alkyl)O-, -OH, (C<sub>1</sub>-C<sub>6</sub>  
 alkyl)S(O)<sub>m</sub>-, (C<sub>1</sub>-C<sub>6</sub> alkyl)C(O)NH-, H<sub>2</sub>N-C(NH)-, (C<sub>1</sub>-C<sub>6</sub>  
 alkyl)C(O)-, (C<sub>1</sub>-C<sub>6</sub> alkyl)OC(O)-, N<sub>3</sub>, (C<sub>1</sub>-C<sub>6</sub> alkyl)OC(O)NH- and  
 20 C<sub>1</sub>-C<sub>20</sub> alkyl.

When R<sup>3a</sup> and R<sup>3b</sup> are combined to form - (CH<sub>2</sub>)<sub>s</sub> -, cyclic  
 moieties are formed. Examples of such cyclic moieties include, but are  
 not limited to:



- 39 -

In addition, such cyclic moieties may optionally include a heteroatom(s). Examples of such heteroatom-containing cyclic moieties include, but are not limited to:



5

The pharmaceutically acceptable salts of the compounds of this invention include the conventional non-toxic salts of the compounds of this invention as formed, e.g., from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those  
 10 derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like: and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pantoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxy-benzoic,  
 15 fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, trifluoroacetic and the like.

It is intended that the definition of any substituent or variable (e.g.,  $R^{10}$ , Z, n, etc.) at a particular location in a molecule be independent of its definitions elsewhere in that molecule. Thus,  
 20  $-N(R^{10})_2$  represents  $-NH_2$ ,  $-NHCH_3$ ,  $-NHC_2H_5$ , etc. It is understood that substituents and substitution patterns on the compounds of the instant invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art as well as those methods set forth below.

25 The pharmaceutically acceptable salts of the compounds of this invention can be synthesized from the compounds of this invention

- 40 -

which contain a basic moiety by conventional chemical methods. Generally, the salts are prepared by reacting the free base with stoichiometric amounts or with an excess of the desired salt-forming inorganic or organic acid in a suitable solvent or various combinations of solvents.

The compounds of the invention can be synthesized from their constituent amino acids by conventional peptide synthesis techniques, and the additional methods described below. Standard methods of peptide synthesis are disclosed, for example, in the following works: Schroeder *et al.*, *"The Peptides"*, Vol. I, Academic Press 1965, or Bodanszky *et al.*, *"Peptide Synthesis"*, Interscience Publishers, 1966, or McOmie (ed.) *"Protective Groups in Organic Chemistry"*, Plenum Press, 1973, or Barany *et al.*, *"The Peptides: Analysis, Synthesis, Biology"* 2, Chapter 1, Academic Press, 1980, or Stewart *et al.*, *"Solid Phase Peptide Synthesis"*, Second Edition, Pierce Chemical Company, 1984. The teachings of these works are hereby incorporated by reference.

Abbreviations used in the description of the chemistry and in the Examples that follow are:

20	Ac <sub>2</sub> O	Acetic anhydride;
	Boc	t-Butoxycarbonyl;
	DBU	1,8-diazabicyclo[5.4.0]undec-7-ene;
	DMAP	4-Dimethylaminopyridine;
	DME	1,2-Dimethoxyethane;
25	DMF	Dimethylformamide;
	EDC	1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide-hydrochloride;
	HOBT	1-Hydroxybenzotriazole hydrate;
	Et <sub>3</sub> N	Triethylamine;
30	EtOAc	Ethyl acetate;
	FAB	Fast atom bombardment;
	HOBT	3-Hydroxy-1,2,2-benzotriazin-4(3H)-one;
	HPLC	High-performance liquid chromatography;
	MCPBA	m-Chloroperoxybenzoic acid;

- 41 -

	MsCl	Methanesulfonyl chloride;
	NaHMDS	Sodium bis(trimethylsilyl)amide;
	Py	Pyridine;
	TFA	Trifluoroacetic acid;
5	THF	Tetrahydrofuran.

Compounds of this invention are prepared by employing the reactions shown in the following Reaction Schemes A-J, in addition to other standard manipulations such as ester hydrolysis, cleavage of  
10 protecting groups, etc., as may be known in the literature or exemplified in the experimental procedures. Some key bond-forming and peptide modifying reactions are:

Reaction A Amide bond formation and subsequent generation of the  
15 amino methyl moiety using standard solution or solid phase methodologies.


Reaction B Preparation of a reduced peptide subunit by reductive  
20 alkylation of an amine by an aldehyde using sodium cyanoborohydride or other reducing agents.

Reaction C Alkylation of the amino moiety of the central phenyl ring.

Reaction D Peptide bond formation and protecting group cleavage using  
25 standard solution or solid phase methodologies.

These reactions may be employed in a linear sequence to provide the compounds of the invention or they may be used to synthesize fragments which are subsequently joined by the alkylation reactions described in the  
30 Reaction Schemes.

- 42 -

REACTION SCHEME AReaction A. Coupling of residues to form an amide bond




- 43 -


REACTION SCHEME B

Reaction B. Preparation of reduced amino acid linkages by reductive alkylation


---



- 44 -

REACTION SCHEME CReaction C. Alkylation of the amino moiety

5

REACTION SCHEME DReaction D. Coupling of residues to form an amide bond10 where  $\text{X}^{\text{L}}$  is a leaving group, e.g.,  $\text{Br}^-$ ,  $\text{I}^-$  or  $\text{MsO}^-$ ;

- 45 -

Reaction Schemes E - M illustrate reactions wherein the non-sulphydryl-containing moiety at the N-terminus of the compounds of the instant invention is attached to an aminomethylbenzamide subunit which may be further elaborated to provide the instant compounds. These reactions may be employed in a linear sequence to provide the compounds of the invention or they may be used to synthesize fragments which are subsequently joined by the reactions described in Reaction Schemes A - D.

The intermediates whose synthesis are illustrated in Reaction Schemes A - D can be reductively alkylated with a variety of aldehydes, such as I, as shown in Reaction Scheme E. The aldehydes can be prepared by standard procedures, such as that described by O. P. Goel, U. Krolls, M. Stier and S. Kesten in Organic Syntheses, 1988, 67, 69-75, from the appropriate amino acid (Reaction Scheme E). The reductive alkylation can be accomplished at pH 5-7 with a variety of reducing agents, such as sodium triacetoxyborohydride or sodium cyanoborohydride in a solvent such as dichloroethane, methanol or dimethylformamide. The product II can be deprotected to give the final compounds III with trifluoroacetic acid in methylene chloride. The final product III is isolated in the salt form, for example, as a trifluoroacetate, hydrochloride or acetate salt, among others. The product diamine III can further be selectively protected to obtain IV, which can subsequently be reductively alkylated with a second aldehyde to obtain V. Removal of the protecting group, and conversion to cyclized products such as the dihydroimidazole VII can be accomplished by literature procedures.

Alternatively, the aminomethylbenzamide subunit can be reductively alkylated with other aldehydes such as 1-trityl-4-carboxaldehyde or 1-trityl-4-imidazolylacetaldehyde, to give products such as VIII (Reaction Scheme F). The trityl protecting group can be removed from VIII to give IX, or alternatively, VIII can first be treated with an alkyl halide then subsequently deprotected to give the alkylated imidazole X. Alternatively, the aminomethylbenzamide subunit can be acylated or sulfonylated by standard techniques.

- 46 -

The imidazole acetic acid **XI** can be converted to the acetate **XIII** by standard procedures, and **XIII** can be first reacted with an alkyl halide, then treated with refluxing methanol to provide the regiospecifically alkylated imidazole acetic acid ester **XIV**. Hydrolysis and reaction with the aminomethylbenzamide subunit in the presence of condensing reagents such as 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (EDC) leads to acylated products such as **XV**.

If the aminomethylbenzamide subunit is reductively alkylated with an aldehyde which also has a protected hydroxyl group, such as **XVI** in Reaction Scheme H, the protecting groups can be subsequently removed to unmask the hydroxyl group (Reaction Schemes H, I). The alcohol can be oxidized under standard conditions to *e.g.* an aldehyde, which can then be reacted with a variety of organometallic reagents such as Grignard reagents, to obtain secondary alcohols such as **XX**. In addition, the fully deprotected amino alcohol **XXI** can be reductively alkylated (under conditions described previously) with a variety of aldehydes to obtain secondary amines, such as **XXII** (Reaction Scheme J), or tertiary amines.

The Boc protected amino alcohol **XVIII** can also be utilized to synthesize 2-aziridinylmethylpiperazines such as **XXIII** (Reaction Scheme K). Treating **XVIII** with 1,1'-sulfonyldiimidazole and sodium hydride in a solvent such as dimethylformamide led to the formation of aziridine **XXIII**. The aziridine reacted in the presence of a nucleophile, such as a thiol, in the presence of base to yield the ring-opened product **XXIV**.


In addition, the aminomethylbenzamide subunit can be reacted with aldehydes derived from amino acids such as O-alkylated tyrosines, according to standard procedures, to obtain compounds such as **XXX**, as shown in Reaction Scheme L. When R' is an aryl group, **XXX** can first be hydrogenated to unmask the phenol, and the amine group deprotected with acid to produce **XXXI**. Alternatively, the amine protecting group in **XXX** can be removed, and O-alkylated phenolic amines such as **XXXII** produced.

- 47 -




Reaction Scheme M illustrates a one pot synthesis of an instant compound wherein the N-terminus nitrogen is substituted with two different non-sulphydryl-containing moieties. Thus, the aminomethylbenzamide subunit is treated with one equivalent of an appropriate aldehyde and, after the reductive adduct has been formed, the in situ intermediate is treated with an equivalent of a different aldehyde.

Similar procedures as are illustrated in Reaction Schemes E-M may be employed using other intermediates such as those whose synthesis is illustrated in Reaction Schemes B - D.


- 48 -

SCHEME E

- 49 -


SCHEME E (continued)

- 50 -


SCHEME F






- 51 -

SCHEME G



- 52 -

SCHEME H


- 53 -

SCHEME I



- 54 -

SCHEME I (CONTINUED)


- 55 -

SCHEME J


- 56 -

SCHEME K

- 57 -




SCHEME L

- 58 -

SCHEME L (continued)




- 59 -

SCHEME M

- 60 -

The  $-NR^4R^5$  moiety of the compounds of the instant invention may provide advantages over a cysteinyl moiety that is incorporated in other types of molecules that are known to be inhibitors of farnesyl protein transferase. In particular, modification of the benzodiazepine compounds described in published PCT application WO 94/26723 with the such  $-NR^4R^5$  substituents as described herein will provide inhibitors of farnesyl protein transferase of the following formulae A and B:



10

A

B

wherein the substituents R, R' and  $R^{25}$  are as defined in WO 94/26723,  $R^{4benz}$ ,  $R^{4'benz}$ ,  $R^{7benz}$  and  $W^{benz}$  are  $R^4$ ,  $R^{4'}$ ,  $R^7$  and W respectively as defined in WO 94/26723 and  $R^a$  and  $R^b$  are defined as  $R^4$  and  $R^5$  are defined herein respectively.


15

- 61 -

Preferably, the following combinations of R<sup>a</sup> and R<sup>b</sup> are selected for incorporation into the compounds of formulae A and B:

R <sup>a</sup>	R <sup>b</sup>
imidazol-4-ylacetyl	H
imidazol-4-ylmethyl	imidazol-4-ylmethyl
4-nitrobenzyl	imidazol-4-ylmethyl
3-(4-cyanobenzyl)-imidazol-4-acetyl	H
3-(4-cyanobenzyl)-imidazol-4-acetyl	CH <sub>3</sub>

- 5 Most preferably, the benzodiazepine compound would be selected from the following formulae:



- wherein R and R' are as defined in WO 94/26723 and W'benz is W' as defined in WO 94/26723 and R<sup>a</sup> and R<sup>b</sup> are defined as R<sup>4</sup> and R<sup>5</sup> are defined herein respectively.
- 10

- Such benzodiazepine analogs may be synthesized by techniques well known in the art, as well as procedures outlined in WO 94/26723. General methods of synthesis of the benzediazapine analogs of this invention are shown in Schemes N, P and Q. Typically a convergent route is employed, which joins the key intermediate 9
- 15

- 62 -

(Scheme N) with suitably functionalized amine and  $R^a$  and  $R^b$  components (Schemes P and Q) using standard amide bond-forming procedures.

- As shown in Scheme N, the protected amino acid **9** may be prepared from a suitably substituted 2-aminobenzophenone (**1**). Many 2-aminobenzophenones are known in the art or are available from commercial sources such as Aldrich Chemical Co. General methods for the synthesis of new 2-aminobenzophenones may be found in the literature (c.f. Walsh, D. A. *Synthesis*, 677-688 (1980).
- 5                   Acylation of **1** with a haloacetyl halide, such as bromoacetyl bromide in a suitable solvent mixture, such as water/  $CH_2Cl_2$ , typically at temperatures ranging from 0° C to 24° C, produces amide **2**. Reaction of **2** with ammonia in a polar solvent such as methanol at 25 to 75 °C then gives the 1,4-benzodiazepin-2-one **3**, after evaporation of the solvent.
- 10                   Alkylation of **3** with a substituted organic ester (**4**), preferably tert-butyl bromacetate, in the presence of a base, preferably  $Cs_2CO_3$  in 1-methyl-2-pyrrolidinone at ambient temperature, gives **5**. Alternatively, **3** may be alkylated at N-1 with a variety of other alkylating agents, for instance, esters of substituted or unsubstituted acrylates, 4-bromobutanoates, etc.
- 15                   Branched compounds (i.e.  $R^{4benz}$  and/or  $R^{4'benz} \neq H$ ), may be prepared by generation of the polyanion of **5** with base and alkylation with an appropriate alkyl halide. Subsequent to alkylation, the ester of **5** may be cleaved with an acid such as TFA (for the tert-butyl esters) or under mild aqueous base hydrolysis (for other alkyl esters) at temperatures between 0
- 20                   and 25 °C.

- 25                   The acid **6** is converted to amino acid **8** via reaction of the dianion, generated with at least two equivalents of a strong base with an electrophilic aminating agent. Alternatively, **6** may be halogenated and reacted with an amine source such as azide (followed by reduction) or ammonia. Preferably, **6** is reacted with 4 equivalents of potassium tert-butoxide in glyme at -5 °C for 30 min and treated with 1.1 equivalents of isobutyl nitrite. The resulting oxime **7** can then be reduced to the racemic amino acid **8** using a variety of reductants, preferably hydrogenation at 40
- 30

- 63 -

psig in the presence of Ruthenium on carbon or Raney nickel in methanol at 50 to 70 °C for 1-4 days.


Amino acid 8 is then suitably protected for selective coupling at the carboxyl terminus. For example, 8 can be converted to the N-BOC derivative 9 using standard amino acid protection conditions, preferably, reaction with equimolar amounts of di-tert-butyl dicarbonate and triethyl amine in DMF/ water at ambient temperature.

For compounds where  $R^a \neq H$ , 9 can be alkylated at nitrogen with a wide variety of alkylating agents including n-alkyl, branched alkyl, and benzyl, according to the standard procedure of Benoiton, et al., *Can. J. Chem.* 1977, 55, 906. For example, reaction of 9 with at least 2 equivalents of base and an alkylating agent in a polar, aprotic solvent at 0 to 50 °C for 0.5 to 48 h give 10. Also, the reactions shown in Schemes E - M may be utilized with the compound 9.


15

- 64 -

**SCHEME N**




- 65 -

SCHEME N(continued)


- Compounds 9 and 10 can be further elaborated according to Schemes P. In general, the carboxylic acid function of 9 and 10 is reacted with a suitably protected amine component using standard solid phase (Scheme P) or solution phase peptide synthesis procedures. The BOC or other protecting group of N-3 of the benzodiazepinone is removed and the amine function then coupled with a third component, for example, a suitably protected amino acid, and then deprotected, again employing standard procedures. The resulting product is subsequently purified by chromatography or crystallization.

- 66 -

SCHEME P




- 67 -

SCHEME P (continued)

- 5                      Alternatively, **3** may be directly alkylated with the "top" sidechain in one intact piece, as shown in Scheme Q. Reaction of **3** with an alkyl halide such as a suitably substituted benzyl bromide, alkyl bromide, in the presence of a base, preferably NaH or Cs<sub>2</sub>CO<sub>3</sub>, gives **11**, which may be processed according to the reactions illustrated in Scheme
- 10    I to provide the desired FPTase inhibitors.

- 68 -

SCHEME Q

- 5 wherein P.G. is a suitably selected protecting group which is utilized if necessary.

- 69 -

The compounds of this invention inhibit Ras farnesyl transferase which catalyzes the first step in the post-translational processing of Ras and the biosynthesis of functional Ras protein. These compounds are useful as pharmaceutical agents for mammals, especially for humans. These compounds may be administered to patients for use in the treatment of cancer. Examples of the type of cancer which may be treated with the compounds of this invention include, but are not limited to, colorectal carcinoma, exocrine pancreatic carcinoma, and myeloid leukemias.

The compounds of this invention may be administered to mammals, preferably humans, either alone or, preferably, in combination with pharmaceutically acceptable carriers or diluents, optionally with known adjuvants, such as alum, in a pharmaceutical composition, according to standard pharmaceutical practice. The compounds can be administered orally or parenterally, including the intravenous, intramuscular, intraperitoneal, subcutaneous, rectal and topical routes of administration.

For oral use of a chemotherapeutic compound according to this invention, the selected compound may be administered, for example, in the form of tablets or capsules, or as an aqueous solution or suspension. In the case of tablets for oral use, carriers which are commonly used include lactose and corn starch, and lubricating agents, such as magnesium stearate, are commonly added. For oral administration in capsule form, useful diluents include lactose and dried corn starch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening and/or flavoring agents may be added. For intramuscular, intraperitoneal, subcutaneous and intravenous use, sterile solutions of the active ingredient are usually prepared, and the pH of the solutions should be suitably adjusted and buffered. For intravenous use, the total concentration of solutes should be controlled in order to render the preparation isotonic.

The present invention also encompasses a pharmaceutical composition useful in the treatment of cancer, comprising the

- 70 -

administration of a therapeutically effective amount of the compounds of this invention, with or without pharmaceutically acceptable carriers or diluents. Suitable compositions of this invention include aqueous solutions comprising compounds of this invention and pharmacologically acceptable carriers, e.g., saline, at a pH level, e.g., 7.4. The solutions may be introduced into a patient's intramuscular blood-stream by local bolus injection.

When a compound according to this invention is administered into a human subject, the daily dosage will normally be determined by the prescribing physician with the dosage generally varying according to the age, weight, and response of the individual patient, as well as the severity of the patient's symptoms.

In one exemplary application, a suitable amount of compound is administered to a mammal undergoing treatment for cancer. Administration occurs in an amount between about 0.1 mg/kg of body weight to about 20 mg/kg of body weight per day, preferably of between 0.5 mg/kg of body weight to about 10 mg/kg of body weight per day.

The compounds of the instant invention are also useful as a component in an assay to rapidly determine the presence and quantity of farnesyl-protein transferase (FPTase) in a composition. Thus the composition to be tested may be divided and the two portions contacted with mixtures which comprise a known substrate of FPTase (for example a tetrapeptide having a cysteine at the amine terminus) and farnesyl pyrophosphate and, in one of the mixtures, a compound of the instant invention. After the assay mixtures are incubated for an sufficient period of time, well known in the art, to allow the FPTase to farnesylate the substrate, the chemical content of the assay mixtures may be determined by well known immunological, radiochemical or chromatographic techniques. Because the compounds of the instant invention are selective inhibitors of FPTase, absence or quantitative reduction of the amount of substrate in the assay mixture without the compound of the instant invention relative to the presence of the unchanged substrate in the assay containing the

- 71 -

instant compound is indicative of the presence of FPTase in the composition to be tested.

It would be readily apparent to one of ordinary skill in the art that such an assay as described above would be useful in identifying tissue samples which contain farnesyl-protein transferase and quantitating the enzyme. Thus, potent inhibitor compounds of the instant invention may be used in an active site titration assay to determine the quantity of enzyme in the sample. A series of samples composed of aliquots of a tissue extract containing an unknown amount of farnesyl-protein transferase, an excess amount of a known substrate of FPTase (for example a tetrapeptide having a cysteine at the amine terminus) and farnesyl pyrophosphate are incubated for an appropriate period of time in the presence of varying concentrations of a compound of the instant invention. The concentration of a sufficiently potent inhibitor (i.e., one that has a  $K_i$  substantially smaller than the concentration of enzyme in the assay vessel) required to inhibit the enzymatic activity of the sample by 50% is approximately equal to half of the concentration of the enzyme in that particular sample.

20

### EXAMPLES

Examples provided are intended to assist in a further understanding of the invention. Particular materials employed, species and conditions are intended to be further illustrative of the invention and not limitative of the reasonable scope thereof.

The standard workup referred to in the examples refers to solvent extraction and washing the organic solution with 10% citric acid, 10% sodium bicarbonate and brine as appropriate. Solutions were dried over sodium sulfate and evaporated *in vacuo* on a rotary evaporator.

30

- 72 -

EXAMPLE 1N-(1(S)-carboxy-3-methylthiopropyl)-3-(4-imidazolylmethyl)-aminomethylbenzamide dihydrochloride

5

Step A: Preparation of N-(1(S)-carbomethoxy-3-methylthiopropyl)-3-chloromethylbenzamide

To a solution of (S) methionine methyl ester hydrochloride (10.56 g, 52.9 mmol) and 4-N-methylmorpholine (21.34 g, 211.6 mmol) under nitrogen in 200 mL of methylene chloride at 0°C was added 3-chloro-methyl benzoyl chloride (10.00 g, 52.9 mmol) dropwise via syringe. After addition the cooling bath was removed and the resulting solution was stirred for 16 h at 20°C. The methylene chloride solution was extracted with 125 mL each of water, 2% potassium hydrogen sulfate, saturated sodium hydrogen carbonate, and saturated sodium chloride. The methylene chloride was dried over magnesium sulfate and concentrated *in vacuo* to the title compound as an oil. <sup>1</sup>HNMR (300 MHz, CDCl<sub>3</sub>) δ 7.85 (1H, s), 7.76 (1H, d, J=8 Hz), 7.56 (1H, d, J=8 Hz), 7.45 (1H, t, J=8 Hz), 6.96 (1H, d, J=7 Hz), 4.94 (1H, q, J=5 Hz), 4.62 (2H, s), 3.81 (3H, s), 2.60 (2H, t, J= 8Hz), 2.30 (1H, m), 2.15 (1H, m), 2.12 (3H, s).

20

Step B: Preparation of N-(1(S)-carbomethoxy-3-methylthiopropyl)-3-azidomethylbenzamide

25

To a stirred solution of the product from Step A (13.52 g, 42.80 mmol) in 50 mL of dimethylsulfoxide under nitrogen was added lithium azide (2.3 g, 47.10 mmol). The solution was stirred for 2 h. The reaction mixture was then partitioned with 300 mL of ethyl acetate and 200 mL of water. The ethyl acetate layer was washed with 125 mL of saturated sodium chloride, dried over magnesium sulfate and concentrated *in vacuo* to afford the title compound as an oil. <sup>1</sup>HNMR (300 MHz, CDCl<sub>3</sub>) δ 7.75 (2H, m), 7.47 (2H, m), 7.02 (1H, d, J=8 Hz), 4.94 (1H, q, J=5 Hz), 4.41 (2H, s), 3.80 (3H, s), 2.60 (2H, t, J=6 Hz), 2.30 (1H, m), 2.15 (1H, m), 2.11 (3H, s).

30

- 73 -

**Step C:**      Preparation of N-(1(S)-carbomethoxy-3-methylthiopropyl)-3-aminomethylbenzamide

To a solution of the product from Step B (11.8 g, 35.08 mmol) in 150 mL of methanol under nitrogen was added 1.5 g 10% palladium on carbon. Hydrogen was applied to the mixture at 1 atmosphere for 1.5 h. The reaction mixture was filtered and concentrated *in vacuo* to obtain 10.3 g (34.76 mmol) of crude product as an oil. The crude product was chromatographed on 500 g of silica gel with chloroform/methanol 95/5 as eluant to afford the title compound as an oil. <sup>1</sup>HNMR (300 MHz, CDCl<sub>3</sub>) δ 7.78 (1H, s), 7.68 (1H, d, J=7 Hz), 7.47 (1H, d, J=7 Hz), 7.40 (1H, t, J=8 Hz), 7.02 (1H, d, J=7 Hz), 4.93 (1H, q, J=5 Hz), 3.92 (2H, s), 3.79 (3H, s), 2.59 (2H, t, J=8 Hz), 2.24 (1H, m), 2.12 (1H, m), 2.10 (3H, s), 1.85 (2H, s).

**Step D:**      Preparation of N-(1(S)-carbomethoxy-3-methylthiopropyl) 3-[(1-triphenylmethyl)-4-imidazolylmethyl]aminomethyl benzamide

To a solution of the product from Step C (0.228 g, 0.767 mmol) in 10 mL of 1,2-dichloroethane was added glacial acetic acid dropwise until a pH 5.5 was achieved. To this mixture at 20°C. was added 0.5 g of crushed 4Å molecular sieves, sodium triacetoxyborohydride (0.487 g, 2.30 mmol), and 1-(triphenylmethyl)-4-imidazole carboxaldehyde (0.130 g, 0.384 mmol). The resulting solution was stirred for 12-72 h. The reaction mixture was filtered through celite and partitioned with 125 mL of water and 150 mL of ethyl acetate. The ethyl acetate layer was washed with 125 mL each of saturated sodium hydrogen carbonate and saturated sodium chloride, dried over magnesium sulfate, and concentrated *in vacuo* to yield 0.363 g of crude product. The crude product was chromatographed on silica gel eluting with chloroform/methanol 95/5 to afford the title compound. <sup>1</sup>HNMR (300 MHz, CDCl<sub>3</sub>) δ 7.83 (1H, s), 7.71 (1H, d, J=7 Hz), 7.40 (4H, m), 7.32 (8H, m), 7.25 (1H, s), 7.12 (7H, m), 6.71 (1H, s), 4.93 (1H, q, J=5

- 74 -

Hz), 3.85 (2H, s), 3.77 (3H, s), 3.71 (2H, s), 2.58 (2H, t, J=8 Hz), 2.25 (1H, m), 2.10 (1H, m), 2.09 (3H, s).

5      **Step E:**      Preparation of N-(1(S)-carbomethoxy-3-methylthiopropyl)-  
3-(4-imidazolylmethyl)aminomethylbenzamide  
dihydrochloride

To a solution of the product from Step D (0.220 g, 0.356 mmol) in 10 mL of methylene chloride was added triethylsilane (0.165 g, 1.42 mmol) and 5 mL of trifluoroacetic acid. The solution was stirred for  
10 45 min, evaporated *in vacuo*, and partitioned with hexane and 0.1% trifluoroacetic acid in water:methanol 2:1. The 0.1% trifluoroacetic acid/water-methanol solution was injected directly onto a Delta-Pak (C-18, 100Å, 15 mm, 40mm x 100mm) prep HPLC column. The gradient at  
15 40 mL/min. was 100% 0.1% TFA/water for 5 min. followed by 95% 0.1% TFA/water :5% 0.1% TFA/acetonitrile to 70% 0.1% TFA/water :30.% 0.1% TFA/acetonitrile over a period of 40 min. The pure fractions were pooled, evaporated *in vacuo* to near dryness, and then taken up in 5 mL of water. This water solution was passed through a 1.2 gm. column of Bio-Rad AG 3-X4 chloride anion exchange resin. The resulting  
20 aqueous column eluant was lyophilized overnight to yield the title compound as a solid. <sup>1</sup>HNMR (300 MHz, CD<sub>3</sub>OD) δ 9.04 (1H, s), 8.06 (1H, s), 7.96 (1H, d, J=8 Hz), 7.83 (1H, s), 7.76 (1H, d, J=8 Hz), 7.63 (1H, t, J=8 Hz), 4.81 (1H, q, J=5 Hz), 4.52 (2H, s), 4.42 (2H, s), 3.77 (3H, s), 2.63 (2H, m), 2.22 (1H, m), 2.16 (1H, m), 2.13 (3H, s). FAB  
25 mass spectrum m/e 377 (m+1). Analysis calculated for C<sub>18</sub>H<sub>24</sub>N<sub>4</sub>O<sub>3</sub>S • 3.3 HCl: C, 43.57; H, 5.55; N, 11.29. Found: C, 43.56; H, 5.54; N, 11.82.

30      **Step F:**      Preparation of N-(1(S)-carboxy-3-methylthiopropyl)-3-(4-imidazolylmethyl)aminomethyl benzamide dihydrochloride

The product from Step E (0.030 g, 0.067mmol) was dissolved in 5 mL of methanol and 3 mL of 5% sodium hydroxide and stirred for 1 h under nitrogen. The reaction mixture was injected directly onto a preparative reverse phase HPLC column with conditions identical



- 75 -

as in the preparation of the compound in Step E. Pure fractions were pooled, evaporated *in vacuo*, and the sample was converted to the hydrochloride salt as before. Lyophilization overnight afforded 0.022 g (0.051 mmol) of the title compound as a solid. <sup>1</sup>HNMR (300 MHz, CD<sub>3</sub>OD) δ 9.06 (1H, s), 8.06 (1H, s), 7.96 (1H, d, J=8 Hz), 7.83 (1H, s), 7.76 (1H, d, J=8 Hz), 7.61 (1H, t, J=8 Hz), 4.78 (1H, q, J=5 Hz), 4.53 (2H, s), 4.47 (2H, s), 2.63 (2H, m), 2.25 (1H, m), 2.15 (1H, m), 2.13 (3H, s). FAB mass spectrum m/e 363 (m+1). Analysis calculated for C<sub>17</sub>H<sub>22</sub>N<sub>4</sub>O<sub>3</sub>S • 3.3 HCl • 0.5 H<sub>2</sub>O: C, 41.57; H, 5.40; N, 11.41. Found: C, 41.54; H, 5.42; N, 11.05.

## EXAMPLE 2

N-(1(S)-carboxy-3-methylthiopropyl)-3-[N,N-bis(4-imidazole-methyl)aminomethyl]benzamide dihydrochloride

Step A: Preparation of N-(1(S)-carbomethoxy-3-methylthiopropyl)-3-[N,N-bis[(1-triphenylmethyl)-4-imidazolylmethyl]-aminomethyl-benzamide

To a solution of the product from Example 1, Step C (0.100 g, 0.357 mmol) in 10 mL of 1,2-dichloroethane was added glacial acetic acid dropwise until the pH was 5.5. To this mixture at 20°C was added 0.5 g of crushed 4Å sieves, sodium triacetoxyborohydride (0.226 g, 2.30 mmol), and 1-(triphenylmethyl)-4-imidazole carboxaldehyde (0.130 g, 1.07 mmol). The resulting solution was stirred for 12-72 h. The reaction mixture was filtered through celite and partitioned with 125 mL of water and 150 mL of ethyl acetate. The ethyl acetate layer was washed with 125 mL each of saturated sodium hydrogen carbonate and saturated sodium chloride, dried over magnesium sulfate, and concentrated *in vacuo* to yield the title compound.

Step B: Preparation of N-(1(S)-carbomethoxy-3-methylthiopropyl)-3-[N,N-bis(4-imidazolylmethyl)aminomethyl]benzamide dihydrochloride

- 76 -

To a solution of the product from Step A (0.320 g, 0.341 mmol) in 10 mL of methylene chloride was added triethylsilane (0.159 g, 1.36 mmol) and 5 mL of trifluoroacetic acid. The solution was stirred for 45 min, evaporated, and partitioned between hexane and 0.1% TFA in water-methanol 2:1. The 0.1% TFA/water:methanol solution was injected directly onto a Delta-Pak (C-18, 100Å, 15 mm, 40mm x 100mm) preparative HPLC column. The gradient at 40 mL/min was 100% 0.1% TFA/water for 5 min followed by 95% 0.1% TFA/water to 60% 0.1% TFA/water :40% 0.1% TFA/acetonitrile over 40 min. The pure fractions were pooled, evaporated to near dryness, and then taken up in 5 mL of water. The aqueous solution was passed through a 1.2 gm. column of Bio-Rad AG 3-X4 chloride anion exchange resin. The resulting aqueous column eluant was lyophilized overnight to yield the title compound as a solid. <sup>1</sup>HNMR (300 MHz, CD<sub>3</sub>OD) δ 8.92 (2H, s), 7.94 (1H, s), 7.78 (1H, d, J=8 Hz), 7.62 (2H, s), 7.58 (1H, d, J=8 Hz), 7.44 (1H, t, J=8 Hz), 4.81 (1H, q, J=5 Hz), 4.06 (4H, s), 3.93 (2H, s), 3.77 (3H, s), 2.63 (2H, m), 2.22 (1H, m), 2.16 (1H, m), 2.13 (3H, s). FAB mass spectrum m/e 457 (m+1). Analysis calculated for C<sub>22</sub>H<sub>28</sub>N<sub>6</sub>O<sub>3</sub>S • 4.8 HCl • 0.2 H<sub>2</sub>O: C, 41.66; H, 5.28; N, 13.25. Found: C, 41.62; H, 5.27; N, 13.02

Step C: Preparation of N-(1(S)-carboxy-3-methylthiopropyl)-3-[N,N-bis(4-imidazolemethyl)aminomethyl]benzamide dihydrochloride

The compound from Step B (0.035 g, 0.052 mmol) was dissolved in 5 mL of methanol and 3 mL of 5% sodium hydroxide and stirred for 1 hr under nitrogen. The reaction mixture was injected directly onto a preparative HPLC column with conditions identical as in Step B. Pure fractions were pooled, evaporated, and the sample converted to the hydrochloride salt as before. Lyophilization overnight afforded the title compound as a solid. <sup>1</sup>HNMR (300 MHz, CD<sub>3</sub>OD) δ 8.88 (2H, s), 7.87 (1H, s), 7.75 (1H, d, J=8 Hz), 7.55 (1H, s), 7.50 (1H, d, J=8 Hz), 7.42 (1H, t, J=8 Hz), 4.78 (1H, q, J=5 Hz), 3.88 (4H, s), 3.77 (2H, s), 2.63 (2H, m), 2.25 (1H, m), 2.15 (1H, m), 2.13 (3H, s). FAB mass spectrum

- 77 -

m/e 443 (m+1). Analysis calculated for  $C_{21}H_{26}N_6O_3S \cdot 5.4 HCl \cdot 1.5 H_2O$ : C, 37.91; H, 5.21; N, 12.63. Found: C, 37.97; H, 5.22; N, 12.37.

### EXAMPLE 3

5

N-(1(S)-carboxy-3-methylthiopropyl) 3-[(4-imidazolylmethyl) -N-methylaminomethyl]benzamide dihydrochloride

10 Step A: Preparation of Methyl 3-chloromethyl benzoate  
To a solution of triethylamine (11.0 mL) in methanol (150 mL) at 0°C was added 3-chloromethylbenzoyl chloride (5.0 g) dropwise. After stirring at 20° C for 0.5 h the solution was concentrated *in vacuo*. The residue was partitioned with 125 mL of water and 150 mL of ethyl acetate. The ethyl acetate layer was washed with 125 mL each of  
15 saturated sodium hydrogen carbonate, 2% potassium hydrogen sulfate and saturated sodium chloride, dried over magnesium sulfate, and concentrated *in vacuo* to yield the title compound.  $^1H$ NMR (300 MHz,  $CDCl_3$ )  $\delta$  8.07 (1H, s), 7.99 (1H, d, J=8 Hz), 7.59 (1H, d, J=8 Hz), 7.43 (1H, t, J=8 Hz), 4.62 (2H, s), 3.92 (3H, s).

20

Step B: Preparation of Methyl 3-azidomethylbenzoate  
Starting with the product from Step A the method used in Step B of Example 1 was used to prepare the title compound.

25

Step C: Preparation of Methyl 3-aminomethyl benzoate  
Starting with the product from Step B the method used in Step C of Example 1 was used to prepare the title compound.

30

Step D: Preparation of Methyl 3-(*t*-butyloxycarbonyl)aminomethyl benzoate

To a solution of the product from Step C (1.14 g) in methylene chloride (50 mL) was added triethylamine (2.90 mL) and di-

- 78 -

tert-butyl dicarbonate (1.67 g) and the mixture was stirred 16 h. The solution was partitioned with water and methylene chloride. The methylene chloride layer was washed with saturated sodium hydrogen carbonate, 2% potassium hydrogen sulfate and saturated sodium chloride, dried over magnesium sulfate, and concentrated *in vacuo* to yield 1.71 g of the crude product. Chromotography on silica gel with hexane/ethyl acetate 9/1 yielded the title compound. <sup>1</sup>HNMR (300 MHz, CDCl<sub>3</sub>) δ 7.95 (1H, s), 7.93 (1H, d, J=8 Hz), 7.49 (1H, d, J=8 Hz), 7.41 (1H, t, J= 8 Hz), 4.90 (1H, b), 4.37 (2H, d, J=6 Hz), 3.92 (3H, s), 1.45 (9H, s).

10

**Step E:** Preparation of Methyl 3- [(t-butyloxycarbonyl)-N-methyl-aminomethyl] benzoate

To a solution of the product from Step E (1.42 g) in dimethylformamide (30 mL) at 0° C was added sodium hydride (0.43 g, 60% dispersion in mineral oil). After stirring for 0.5 h methyl iodide (0.40 mL) was added and the mixture was stirred 16 h at 20° C. The solution was concentrated *in vacuo* and the residue was partitioned between ethyl acetate and water. The ethyl acetate layer was washed with saturated sodium hydrogen carbonate, 2% potassium hydrogen sulfate and saturated sodium chloride, dried over magnesium sulfate, and concentrated *in vacuo* to yield the crude product. Chromotography on silica gel with hexane/ethyl acetate 9/1 yielded 0.35 g of the title compound. <sup>1</sup>HNMR (300 MHz, CDCl<sub>3</sub>) δ 7.93 (2H, m), 7.42 (2H, m), 4.45 (2H, s), 3.92 (3H, s), 2.83 (3H, d), 1.47 (9H, s).

25

**Step F:** Preparation of 3- [(t-butyloxycarbonyl)-N-methyl-aminomethyl] benzoic acid

To a solution of the product from Step E (0.35 g) in methanol was added 5% sodium hydroxide. After stirring for 2 h the methanol was evaporated and the aqueous layer was adjusted to pH 3 with 2% potassium hydrogen sulfate. The aqueous layer was extracted with ethyl acetate several times. The ethyl acetate layer was washed with saturated sodium chloride, dried over magnesium sulfate, and concentrated *in vacuo* to yield the title compound.

30

- 79 -

**Step G:** Preparation of N-(1(S)-carbomethoxy-3-methylthiopropyl)-  
3-[(t-butyloxycarbonyl)-N-methylaminomethyl]benzamide

To a solution of the product from Step F (0.27 g) in  
5 dimethylformamide (10 mL) was added hydroxybenzotriazole (0.16 g),  
EDC (0.19 g), N-methylmorpholine (0.40 mL), and (S) methionine methyl  
ester hydrochloride (0.203 mg). After stirring for 2 h the solution was  
concentrated *in vacuo* and the residue was partitioned with water and  
ethyl acetate. The ethyl acetate layer was washed with saturated sodium  
10 hydrogen carbonate, 2% potassium hydrogen sulfate and saturated  
sodium chloride, dried over magnesium sulfate, and concentrated *in vacuo*  
to yield the title compound. <sup>1</sup>HNMR (300 MHz, CDCl<sub>3</sub>) δ 7.70  
(2H, s), 7.40 (2H, m), 6.95 (1H, d, J=7 Hz), 4.94 (1H, q, J=7 Hz), 4.45  
(2H, s), 3.81 (3H, s), 2.82 (3H, d), 2.59 (2H, m), 2.30 (1H, m), 2.13 (1H,  
15 m), 2.12 (3H, s), 1.46 (9H, s).

**Step H:** Preparation of N-(1(S)-carbomethoxy-3-methylthiopropyl)-  
3-(N-methylaminomethyl)benzamide trifluoroacetate

To a solution of the product from Step G in methylene  
20 chloride was added trifluoroacetic acid (33% by volume). After stirring  
for 1 h the solution was concentrated *in vacuo* to yield the title  
compound.

**Step I:** Preparation of N-(1(S)-carbomethoxy-3-methylthiopropyl)-  
25 3-[(1-triphenylmethyl)-4-imidazolylmethyl-N-methyl-  
aminomethyl]benzamide

Starting with the product from Step H (0.18 g) the method  
described in Step D of Example 1 was used to prepare the title  
compound.

30

**Step J:** Preparation of N-(1(S)-carbomethoxy-3-methylthiopropyl)-  
3-[(4-imidazolylmethyl)-N-methyl-minomethyl]benzamide  
dihydrochloride

- 80 -

Starting with the compound from Step I (0.24 g) the method described in Step E of Example 1 was used to prepare the title compound. FAB mas spectrum m/e 391 (m+1).

Analysis for  $C_{19}H_{26}N_4O_3S \cdot 5.0 HCl \cdot 0.5 H_2O$ :

5    Calculated: C, 39.32; H, 5.56; N, 9.65;

Found:        C, 39.33; H, 5.57; N, 9.38.

10    Step K:        Preparation of N-(1(S)-carboxy-3-methylthiopropyl)-3-[(4-imidazolomethyl) -N-methyl-aminomethyl]benzamide dihydrochloride

Starting with the compound from Step J (0.035 g) the method described in Step F of Example 1 was used to prepare the above title compound. FAB mas spectrum m/e 377 (m+1).

Analysis for  $C_{18}H_{24}N_4O_3S \cdot 3.70 HCl \cdot 0.2 H_2O$ :

15    Calculated    C, 42.05; H, 5.51; N, 10.90;

Found:        C, 42.09; H, 5.49; N, 10.70.

#### EXAMPLE 4

20    N-(1(S)-carboxy-3-methylthiopropyl) 4-[(4-imidazolylmethyl)-amino]benzamide dihydrochloride

Step A:        Preparation of 4-t-butyloxycarbonyl- aminobenzoic acid

25    Starting with 4-aminobenzoic acid (2.00 g) dissolved in tetrahydrofuran (50 mL) and 5% sodium hydroxide (15 mL) the method described in Step D of Example 3 was used to prepare the title compound. After extractive work up obtained the title compound.

$^1H$ NMR (300 MHz,  $CDCl_3$ )  $\delta$  8.04 (2H, d, J=9 Hz), 7.46 (2H, d, J=9 Hz), 6.75 (1H, s), 1.46 (9H, s).

30

Step B:        Preparation of N-(1(S)-carbomethoxy-3-methylthiopropyl)-4-(t-butyloxycarbonyl)aminobenzamide

To a solution of the product from Step A (0.5 g) in dimethylformamide (20 mL) was added hydroxybenzotriazole (0.37 g),

- 81 -

EDC (0.51 g), N-methylmorpholine 0.8 mL), and (S) methionine methyl ester hydrochloride (0.49 g). After stirring for 16 h the solution was concentrated *in vacuo* and the residue was partitioned with water and ethyl acetate. The ethyl acetate layer was washed with saturated sodium hydrogen carbonate, 2% potassium hydrogen sulfate and saturated sodium chloride, dried over magnesium sulfate, and concentrated *in vacuo* to yield the title compound.

10 Step C: Preparation of N-(1(S)-carbomethoxy-3-methylthiopropyl)-4-aminobenzamide

To a solution of the product from Step B in methylene chloride was added trifluoroacetic acid (33% by volume). After stirring for 1 h the solution was concentrated *in vacuo* to yield the trifluoroacetate salt of the product (0.59 g). This product was partitioned between ethyl acetate and saturated sodium bicarbonate. The ethyl acetate layer was washed with saturated sodium chloride, dried over magnesium sulfate, and concentrated *in vacuo* to yield the title compound. <sup>1</sup>HNMR (300 MHz, CDCl<sub>3</sub>) δ 7.64 (2H, d, J=8 Hz), 6.75 (1H, d, J=8 Hz), 6.65 (2H, d, J=8 Hz), 4.90 (1H, q, J=5 Hz), 3.78 (3H, s), 2.57 (2H, m), 2.27 (1H, m), 2.08 (4H, m).

25 Step D: Preparation of N-(1(S)-carbomethoxy-3-methylthiopropyl)-4-[(1-triphenylmethyl)-4-imidazolylmethyl]-aminobenzamide

Starting with the product from Step C (0.07 g) the method described in Step D of Example 1 was used to prepare the title compound.

30 Step E: Preparation of N-(1(S)-carbomethoxy-3-methylthiopropyl)-4-(4-imidazolylmethyl)aminobenzamide dihydrochloride

Starting with the product from Step D (0.24 g) the method described in Step E of Example 1 was used to prepare the title compound. FAB mas spectrum m/e 363 (m+1). Analysis for C<sub>17</sub>H<sub>22</sub>N<sub>4</sub>O<sub>3</sub>S • 2.8 HCl:

- 82 -

Calculated: C, 43.99; H, 5.39; N, 12.07;

Found: C, 43.94; H, 5.37; N, 12.24.

5     **Step F:**     Preparation of N-(1(S)-carboxy-3-methylthiopropyl)-4-[(4-  
                  imidazolylmethyl)amino]benzamide dihydrochloride  
                  Starting with (5) (0.035 g) the method described in Step F of  
Example 1 was used to prepare the title compound. FAB mas spectrum  
m/e 349 (m+1).

Analysis for  $C_{16}H_{20}N_4O_3S \cdot 3.10 \text{ HCl} \cdot 1.2 \text{ H}_2\text{O}$ :

10    Calculated: C, 39.91; H, 5.13; N, 11.63;

Found: C, 39.87; H, 5.12; N, 11.22.

Using the appropriate starting materials the methods described above for  
Example 4 were used to prepare Examples 5-7.

15

### EXAMPLE 5

N-(1(S)-carboxy-3-methylthiopropyl)-3-[(4-imidazolylmethyl) amino]  
benzamide dihydrochloride

20

FAB mass spectrum m/e 349 (m+1).

Analysis for  $C_{16}H_{20}N_4O_3S \cdot 3.2 \text{ HCl}$ :

Calculated: C, 41.37; H, 5.03; N, 12.06;

Found: C, 41.32; H, 4.92; N, 11.69.

25



- 83 -

EXAMPLE 6

N-(1(S)-carbomethoxy-3-methylthiopropyl)-3-[(4-imidazolylpropyl) amino]benzamide dihydrochloride

---

5

FAB mass spectrum m/e 391 (m+1).

Analysis for  $C_{19}H_{26}N_4O_3S \cdot 2.4 HCl \cdot 1.3 H_2O$ :

Calculated: C, 45.53; H, 6.23; N, 11.18;

Found: C, 45.51; H, 6.25; N, 11.10.

10

EXAMPLE 7

N-(1(S)-Carboxy-3-methylthiopropyl)-3-[(4-imidazolylpropyl) amino]benzamide dihydrochloride

---

15

FAB mass spectrum m/e 377 (m+1).

Analysis for  $C_{18}H_{24}N_4O_3S \cdot 3.0 HCl \cdot 0.5 H_2O$ :

Calculated: C, 43.73; H, 5.71; N, 11.33;

Found: C, 43.67; H, 5.71; N, 10.72.

20

EXAMPLE 8

N-(1(S)-Carboxy-3-methylthiopropyl)-3-[N-(4-imidazolylmethyl)-N-(4-nitrobenzyl)aminomethyl]benzamide ditrifluoroacetate

---

25

Step A: N-(1(S)-Carbomethoxy-3-methylthiopropyl)-3-[N-(4-imidazolylmethyl)-N-(4-nitrobenzyl)aminomethyl]-benzamide ditrifluoroacetate

---

30 N-(1(S)-carbomethoxy-3-methylthiopropyl)-3-

aminomethylbenzamide (0.104 g, 0.352 mmol) was dissolved in dichloroethane (5 mL). Crushed molecular sieves (0.209 g) and sodium triacetoxyborohydride (0.186 g, 0.881 mmol). The pH was about 7.5. 4-Nitrobenzaldehyde (0.0533 g, 0.352 mmol) was added plus 0.5 drop of acetic acid to bring the pH to about 7. The reaction was stirred 2 h under

- 84 -

nitrogen at 20°C. 1-Triphenylmethylimidazolyly-4-carboxaldehyde (0.119 g, 0.352 mmol) was added to the reaction mixture with additional sodium triacetoxyborohydride and dichloroethane (2 mL). Triethylamine (5 drops) brought the pH to about 7. The reaction continued to stir at 20°C under nitrogen overnight. The reaction was quenched with saturated sodium bicarbonate solution and let stir 20 min. It was then removed to a separatory funnel with copious amounts of ethyl acetate. The aqueous layer was removed and the organic phase was washed with saturated brine and dried over magnesium sulfate. The crude product was chromatographed on silica gel with 50% ethyl acetate in hexane. This chromatographed product was dissolved in dichloromethane (7 mL); triethylsilane (0.5 mL, 3.13 mmol) was added and then trifluoroacetic acid (3.5 mL). After 0.5 h at 20°C, the solvent was evaporated and the residue partitioned between hexane and water. The aqueous solution was purified by preparative reverse phase HPLC using a 100 mm Waters PrepPak® reverse phase column (DeltaPak™ C18, 50µM, 100 Å) and pure product isolated by gradient elution using 80% 0.1% trifluoroacetic acid in water (Solvent A) and 20% 0.1% trifluoroacetic acid in acetonitrile (Solvent B) to 55% Solvent A and 45% Solvent B. The pure fractions were combined and the solvent evaporated, and the pure product was dissolved in water and lyophilized to give the title compound as a clear, pale yellow solid. <sup>1</sup>HNMR (CD<sub>3</sub>OD, 400 MHz) δ 8.78 (1H, br s), 8.18 (2H, d, J=8.6 Hz), 7.86 (1H br s), 7.74 (1H, br d, J=8 Hz), 7.64 (2H, d, J=8.6 Hz), 7.56 (1H, br d, J=8 Hz), 7.46 (1H, br s), 7.44 (1H, dd J=8, 8 Hz), 4.8 (1H, m), 3.74 to 3.79 (9H, m), 2.58 to 2.66 (2H, m), 2.23 (1H, m), 2.12 (1H, m), 2.10 (3H, s). FAB ms (m+1) 512. Anal. Calc. for C<sub>25</sub>H<sub>29</sub>N<sub>5</sub>O<sub>5</sub>S · 0.70 H<sub>2</sub>O · 3.30 TFA. Found: C, 42.12; H, 3.75, N, 7.91.

30 Step B: Preparation of N-(1(S)-carboxy-3-methylthiopropyl)-3-[N-(4-imidazolylylmethyl)-N-(4-nitrobenzyl)aminomethyl]-benzamide ditrifluoroacetate

The product from Step A (0.045 g, 0.0608 mmol) was dissolved in methanol (4 mL) and 0.5 mL of 10% NaOH solution was

- 85 -

added to take pH to about 12. Water (4 mL) was added. At 3 h reaction was purified and lyophilized according to the procedure described in Step A to the title compound as a white solid. <sup>1</sup>HNMR (CD<sub>3</sub>OD, 400 MHz) δ 8.78 (1H, br s), 8.18 (2H, d, J=8.6 Hz), 7.88 (1H, br s), 7.75 (1H, br d, J=8 Hz), 7.65 (2H, d, J=8.6 Hz), 7.55 (1H, br d, J=8 Hz), 7.46 (1H, br s), 7.43 (1H, dd, J=8, 8 Hz), 4.8 (1H, m), 3.80 (4H, br s), 3.75 (2H, br s), 2.58 to 2.68 (2H, m), 2.27 (1H, m), 2.13 (1H, m), 2.11 (3H, s). FAB ms (m+1) 498, Anal Calc. for C<sub>24</sub>H<sub>27</sub>N<sub>5</sub>O<sub>5</sub>S · 1.40 H<sub>2</sub>O + 3.20 TFA. Found: C, 41.16; H, 3.72; N, 8.11.

### EXAMPLE 9

N-(1(S)-carboxy-3-methylthiopropyl)-3-[N,N-bis-(4-nitrophenylmethyl)aminomethyl]benzamide dihydrochloride

Step A: Preparation of N-(1(S)-carboxymethyl-3-methylthiopropyl)-3-[N,N-bis-(4-nitrophenylmethyl)] ditrifluoroacetate

The product from Example 1, Step 3 (0.100 g, 0.337 mmol) was dissolved in dichloroethane (5 mL). *p*-Nitrobenzaldehyde, sodium triacetoxymethylborohydride (0.214 g, 1.01 mmol) and crushed molecular sieves were added, and the pH adjusted to 5.5 with acetic acid and triethylamine. The reaction was stirred at 20°C overnight, quenched with saturated sodium bicarbonate, and partitioned between ethyl acetate and saturated sodium bicarbonate. The organic phase was washed with 2% potassium hydrogen sulfate, saturated sodium bicarbonate, saturated brine, and dried over magnesium sulfate. The crude product was purified by silica gel chromatography using 40% ethyl acetate in hexane. This product was further purified by preparative reverse phase HPLC using a gradient elution from 85% water, 15% acetonitrile to 20% water over a period of 40 min. (solvents contained 0.1% trifluoroacetic acid). <sup>1</sup>HNMR (300 MHz, CDCl<sub>3</sub>) δ 8.25(4H, d, J=8.5 Hz), 7.92 (1H, s), 7.80 (1H, d, J=7.6 Hz), 7.63 (4H, d, J=8.5 Hz), 7.53 (m, 2H), 7.35 (1H, d, J=7.3 Hz), 4.94 (1H, bq, J=6.2 Hz), 3.98 (4H, s), 3.95 (2H, s), 3.82 (3H, s), 2.61 (2H, t, J=7.3 Hz), 2.30 (1H, m), 2.19 (1H, dt, J=15, 7.5 Hz), 2.11

- 86 -

(3H, s). Analysis calculated for  $C_{28}H_{30}N_4O_7S \cdot 2.1 CF_3CO_2H \cdot 0.5 H_2O$ : C, 47.45; H, 4.09; N, 6.87. Found: C, 47.44; H, 4.01; N, 6.91.

5     **Step B:**     Preparation of N-(1(S)-carboxy-3-methylthiopropyl)-3-[N,N-bis-(4-nitrophenylmethyl)aminomethyl]benzamide ditrifluoroacetate

10     The product from Step 1 (0.025 g) was hydrolyzed to the acid according to the procedure described in Example 1, Step 6. The title compound was obtained after purification by preparative reverse phase HPLC. FAB ms m/e (m+1) 553). Analysis calculated for  $C_{27}H_{28}N_4O_7S \cdot 1.6 CF_3CO_2H \cdot 0.2 H_2O$ : C, 49.11; H, 4.09; N, 7.59. Found: C, 49.10; H, 3.93; N, 7.55.

### EXAMPLE 10

15

#### *In vitro* inhibition of ras farnesyl transferase

Assays of farnesyl-protein transferase. Partially purified bovine FPTase and Ras peptides (Ras-CVLS, Ras-CVIM and RAS-CAIL) were prepared as described by Schaber *et al.*, *J. Biol. Chem.* 265:14701-14704 (1990),  
20     Pompliano, *et al.*, *Biochemistry* 31:3800 (1992) and Gibbs *et al.*, *PNAS U.S.A.* 86:6630-6634 (1989), respectively. Bovine FPTase was assayed in a volume of 100  $\mu$ l containing 100 mM N-(2-hydroxy ethyl) piperazine-N'-(2-ethane sulfonic acid) (HEPES), pH 7.4, 5 mM  $MgCl_2$ , 5 mM dithiothreitol (DTT), 100 mM [ $^3H$ ]-farnesyl diphosphate ([ $^3H$ ]-FPP;  
25     740 CBq/mmol, New England Nuclear), 650 nM Ras-CVLS and 10  $\mu$ g/ml FPTase at 31°C for 60 min. Reactions were initiated with FPTase and stopped with 1 ml of 1.0 M HCL in ethanol. Precipitates were collected onto filter-mats using a TomTec Mach II cell harvester, washed with 100% ethanol, dried and counted in an LKB  $\beta$ -plate counter. The  
30     assay was linear with respect to both substrates, FPTase levels and time; less than 10% of the [ $^3H$ ]-FPP was utilized during the reaction period. Purified compounds were dissolved in 100% dimethyl sulfoxide (DMSO) and were diluted 20-fold into the assay. Percentage inhibition is measured by the amount of incorporation of radioactivity in the presence

- 87 -

of the test compound when compared to the amount of incorporation in the absence of the test compound.

Human FPTase was prepared as described by Omer *et al.*, *Biochemistry* 32:5167-5176 (1993). Human FPTase activity was assayed as described above with the exception that 0.1% (w/v) polyethylene glycol 20,000, 10  $\mu$ M ZnCl<sub>2</sub> and 100 nM Ras-CVIM were added to the reaction mixture. Reactions were performed for 30 min., stopped with 100  $\mu$ l of 30% (v/v) trichloroacetic acid (TCA) in ethanol and processed as described above for the bovine enzyme.

The compounds of the instant invention were tested for inhibitory activity against human FPTase by the assay described above and were found to have IC<sub>50</sub> of < 100  $\mu$ M.

### EXAMPLE 11

15

#### In vivo ras farnesylation assay

The cell line used in this assay is a v-ras line derived from either Rat1 or NIH3T3 cells, which expressed viral Ha-ras p21. The assay is performed essentially as described in DeClue, J.E. *et al.*, *Cancer Research* 51:712-717, (1991). Cells in 10 cm dishes at 50-75% confluency are treated with the test compound (final concentration of solvent, methanol or dimethyl sulfoxide, is 0.1%). After 4 hours at 37°C, the cells are labelled in 3 ml methionine-free DMEM supplemented with 10% regular DMEM, 2% fetal bovine serum and 400 mCi[<sup>35</sup>S]methionine (1000 Ci/mmol). After an additional 20 hours, the cells are lysed in 1 ml lysis buffer (1% NP40/20 mM HEPES, pH 7.5/5 mM MgCl<sub>2</sub>/1mM DTT/10 mg/ml aprotinin/2 mg/ml leupeptin/2 mg/ml antipain/0.5 mM PMSF) and the lysates cleared by centrifugation at 100,000 x g for 45 min. Aliquots of lysates containing equal numbers of acid-precipitable counts are brought to 1 ml with IP buffer (lysis buffer lacking DTT) and immunoprecipitated with the ras-specific monoclonal antibody Y13-259 (Furth, M.E. *et al.*, *J. Virol.* 43:294-304, (1982)). Following a 2 hour antibody incubation at 4°C, 200  $\mu$ l of a 25% suspension of protein A-Sepharose coated with rabbit anti rat IgG is

- 88 -

added for 45 min. The immunoprecipitates are washed four times with IP buffer (20 mM HEPES, pH 7.5/1 mM EDTA/1% Triton X-100/0.5% deoxycholate/0.1%/SDS/0.1 M NaCl) boiled in SDS-PAGE sample buffer and loaded on 13% acrylamide gels. When the dye front reached the bottom, the gel is fixed, soaked in Enlightening, dried and autoradiographed. The intensities of the bands corresponding to farnesylated and nonfarnesylated ras proteins are compared to determine the percent inhibition of farnesyl transfer to protein.

10

## EXAMPLE 12

### *In vivo* growth inhibition assay


To determine the biological consequences of FPTase inhibition, the effect of the compounds of the instant invention on the anchorage-independent growth of Rat1 cells transformed with either a v-*ras*, v-*raf*, or v-*mos* oncogene is tested. Cells transformed by v-Raf and v-Mos maybe included in the analysis to evaluate the specificity of instant compounds for Ras-induced cell transformation.

Rat 1 cells transformed with either v-*ras*, v-*raf*, or v-*mos* are seeded at a density of  $1 \times 10^4$  cells per plate (35 mm in diameter) in a 0.3% top agarose layer in medium A (Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum) over a bottom agarose layer (0.6%). Both layers contain 0.1% methanol or an appropriate concentration of the instant compound (dissolved in methanol at 1000 times the final concentration used in the assay). The cells are fed twice weekly with 0.5 ml of medium A containing 0.1% methanol or the concentration of the instant compound. Photomicrographs are taken 16 days after the cultures are seeded and comparisons are made.

- 89 -

WHAT IS CLAIMED IS:

1. A compound which inhibits Ras farnesyl-transferase having the Formula I:



wherein:

R<sup>1a</sup> and R<sup>1b</sup> are independently selected from:

- 10      a) hydrogen,  
          b) aryl, heterocycle, cycloalkyl, alkenyl, alkynyl, R<sup>10</sup>O-,  
                  R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>,  
                  (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>,  
                  or R<sup>11</sup>OC(O)NR<sup>10</sup>-,  
 15      c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl,  
                  heterocyclic, cycloalkyl, alkenyl, alkynyl, R<sup>10</sup>O-,  
                  R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-,  
                  R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)-  
                  NR<sup>10</sup>-;

20      R<sup>2a</sup> and R<sup>2b</sup> are independently selected from:

- a) hydrogen,  
          b) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by alkenyl, R<sup>10</sup>O-,  
                  R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, N<sub>3</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-,  
                  R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-,  
 25      c) aryl, heterocycle, cycloalkyl, alkenyl, R<sup>10</sup>O-,  
                  R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>,  
                  (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>,  
                  -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and

- 90 -

- d) C<sub>1</sub>-C<sub>6</sub> alkyl substituted with an unsubstituted or substituted group selected from aryl, heterocyclic and C<sub>3</sub>-C<sub>10</sub> cycloalkyl;

5 R<sup>3a</sup> and R<sup>3b</sup> are independently selected from:


- a) a side chain of a naturally occurring amino acid,  
 b) an oxidized form of a side chain of a naturally occurring amino acid which is:  
 i) methionine sulfoxide, or  
 10 ii) methionine sulfone, and  
 c) substituted or unsubstituted C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, aryl or heterocyclic group,  
 wherein the substituent is selected from F, Cl, Br, N(R<sup>10</sup>)<sub>2</sub>, NO<sub>2</sub>, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-,  
 15 CN, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, R<sup>11</sup>OC(O)NR<sup>10</sup>- and C<sub>1</sub>-C<sub>20</sub> alkyl, and  
 d) C<sub>1</sub>-C<sub>6</sub> alkyl substituted with an unsubstituted or substituted group selected from aryl, heterocycle and C<sub>3</sub>-C<sub>10</sub> cycloalkyl; or

20

R<sup>3a</sup> and R<sup>3b</sup> are combined to form -(CH<sub>2</sub>)<sub>s</sub> - wherein one of the carbon atoms is optionally replaced by a moiety selected from: O, S(O)<sub>m</sub>, -NC(O)-, and -N(COR<sup>10</sup>)- ;

25 R<sup>4</sup> and R<sup>5</sup> are independently selected from:

- a) hydrogen, and  
 b)



30 R<sup>7</sup> is independently selected from:

- a) hydrogen,



- 91 -

- 5           b) aryl, heterocycle, cycloalkyl, alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $NO_2$ ,  $R^{10}_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ , and
- 10           c)  $C_1-C_6$  alkyl unsubstituted or substituted by aryl, heterocycle, cycloalkyl, alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NH-$ , CN,  $H_2N-C(NH)-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{10}OC(O)NH-$ ;
- 15           R<sup>8</sup> is selected from:
- a) hydrogen,
- b) alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $NO_2$ ,  $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ , and
- 20           c)  $C_1-C_6$  alkyl unsubstituted or substituted by perfluoroalkyl, F, Cl, Br,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ ;

R<sup>10</sup> is independently selected from hydrogen,  $C_1-C_6$  alkyl, benzyl and aryl;

25 R<sup>11</sup> is independently selected from  $C_1-C_6$  alkyl and aryl;

A<sup>1</sup> and A<sup>2</sup> are independently selected from: a bond,  $-CH=CH-$ ,  $-C\equiv C-$ ,  $-C(O)-$ ,  $-C(O)NR^{10}-$ ,  $-NR^{10}C(O)-$ , O,  $-N(R^{10})-$ ,  $-S(O)_2N(R^{10})-$ ,  $-N(R^{10})S(O)_2-$ , or  $S(O)_m$ ;

30

V is selected from:

- a) hydrogen,
- b) heterocycle,
- c) aryl,

- 92 -

- d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
- e) C<sub>2</sub>-C<sub>20</sub> alkenyl,
- provided that V is not hydrogen if A<sup>1</sup> is S(O)<sub>m</sub> and V is not hydrogen if
- 5 A<sup>1</sup> is a bond, n is 0 and A<sup>2</sup> is S(O)<sub>m</sub>;

W is a heterocycle;


Z is independently H<sub>2</sub> or O;

10

- m is 0, 1 or 2;
- n is 0, 1, 2, 3 or 4;
- p is 0, 1, 2, 3 or 4;
- r is 0 to 5, provided that r is 0 when V is hydrogen;
- 15 s is 4 or 5; and
- u is 0 or 1;

or a pharmaceutically acceptable salt thereof.

- 20 2. A prodrug of the compound of Claim 1 having the  
Formula II:



- 25 wherein:
- R<sup>1a</sup> and R<sup>1b</sup> are independently selected from:
- a) hydrogen,

- 93 -

- b) aryl, heterocycle, cycloalkyl, alkenyl, alkynyl,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $NO_2$ ,  $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ ,
- 5 c)  $C_1-C_6$  alkyl unsubstituted or substituted by aryl, heterocyclic, cycloalkyl, alkenyl, alkynyl,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ ;

10

$R^{2a}$  and  $R^{2b}$  are independently selected from:

- a) hydrogen,
- b)  $C_1-C_6$  alkyl unsubstituted or substituted by alkenyl,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $N_3$ ,  $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ ,
- 15 c) aryl, heterocycle, cycloalkyl, alkenyl,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $NO_2$ ,  $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ , and
- 20 d)  $C_1-C_6$  alkyl substituted with an unsubstituted or substituted group selected from aryl, heterocyclic and  $C_3-C_{10}$  cycloalkyl;

$R^{3a}$  and  $R^{3b}$  are independently selected from:

- 25 a) a side chain of a naturally occurring amino acid,
- b) an oxidized form of a side chain of a naturally occurring amino acid which is:
- i) methionine sulfoxide, or
- ii) methionine sulfone, and
- 30 c) substituted or unsubstituted  $C_1-C_{20}$  alkyl,  $C_2-C_{20}$  alkenyl,  $C_3-C_{10}$  cycloalkyl, aryl or heterocyclic group,
- wherein the substituent is selected from F, Cl, Br,  $N(R^{10})_2$ ,  $NO_2$ ,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ ,

- 94 -


CN,  $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  
 $-N(R^{10})_2$ ,  $R^{11}OC(O)NR^{10}-$  and  $C_1-C_{20}$  alkyl, and

- d)  $C_1-C_6$  alkyl substituted with an unsubstituted or substituted group selected from aryl, heterocycle and  $C_3-C_{10}$  cycloalkyl; or

$R^{3a}$  and  $R^{3b}$  are combined to form  $-(CH_2)_s-$  wherein one of the carbon atoms is optionally replaced by a moiety selected from: O,  $S(O)_m$ ,  $-NC(O)-$ , and  $-N(COR^{10})-$ ;

$R^4$  and  $R^5$  are independently selected from:

- a) hydrogen, and  
 b)




$R^6$  is

- a) substituted or unsubstituted  $C_1-C_8$  alkyl or substituted or unsubstituted  $C_5-C_8$  cycloalkyl, wherein the substituent on the alkyl is selected from:

- 1) aryl,  
 2) heterocycle,  
 3)  $-N(R^{11})_2$ ,  
 4)  $-OR^{10}$ , or

b)



$R^7$  is independently selected from:

- a) hydrogen,  
 b) aryl, heterocycle, cycloalkyl, alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,

$R^{10}C(O)NR^{10}$ -,  $CN$ ,  $NO_2$ ,  $R^{10}_2N-C(NR^{10})$ -,  $R^{10}C(O)$ -,  $R^{10}OC(O)$ -,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}$ -, and

5 c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl, heterocycle, cycloalkyl, alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NH-, CN, H<sub>2</sub>N-C(NH)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>10</sup>OC(O)NH-;

**R8 is selected from:**

10            a)    **hydrogen,**

b) alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br, R<sup>10</sup>O-, R<sup>11</sup>S(O)m-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C-(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and

15 c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by perfluoroalkyl, F, Cl, Br, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-;

20 R<sup>10</sup> is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl and aryl;

**R<sup>11</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;**

25 R<sup>12</sup> is independently selected from hydrogen and C<sub>1</sub>-C<sub>6</sub> alkyl;

**R<sup>13</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl;**

30 A<sup>1</sup> and A<sup>2</sup> are independently selected from: a bond, -CH=CH-, -C≡C-,  
-C(O)-, -C(O)NR<sup>10</sup>-, -NR<sup>10</sup>C(O)-, O, -N(R<sup>10</sup>)-,  
-S(O)<sub>2</sub>N(R<sup>10</sup>)-, -N(R<sup>10</sup>)S(O)<sub>2</sub>-, or S(O)<sub>m</sub>;

**V is selected from:**

a) hydrogen,

- 96 -


- b) heterocycle,  
 c) aryl,  
 d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and  
 5 e) C<sub>2</sub>-C<sub>20</sub> alkenyl,  
 provided that V is not hydrogen if A<sup>1</sup> is S(O)<sub>m</sub> and V is not hydrogen if A<sup>1</sup> is a bond, n is 0 and A<sup>2</sup> is S(O)<sub>m</sub>;

W is a heterocycle;  
 10 Z is independently H<sub>2</sub> or O;

m is 0, 1 or 2;  
 n is 0, 1, 2, 3 or 4;  
 15 p is 0, 1, 2, 3 or 4;  
 r is 0 to 5, provided that r is 0 when V is hydrogen;  
 s is 4 or 5; and  
 u is 0 or 1;

20 or a pharmaceutically acceptable salt thereof.

3. A compound which inhibits Ras farnesyl-transferase having the Formula III:



25

wherein:  
 R<sup>1a</sup> and R<sup>1b</sup> are independently selected from:

- 97 -


- 5 a) hydrogen,  
 b) aryl, heterocycle, cycloalkyl, alkenyl, alkynyl,  $R^{10}O-$ ,  
 $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $NO_2$ ,  
 $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ ,  
 or  $R^{11}OC(O)NR^{10}-$ ,  
 10 c)  $C_1-C_6$  alkyl unsubstituted or substituted by aryl,  
 heterocyclic, cycloalkyl, alkenyl, alkynyl,  $R^{10}O-$ ,  
 $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $(R^{10})_2N-C(NR^{10})-$ ,  
 $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)-$   
 15  $NR^{10}-$ ;

$R^{2a}$  and  $R^{2b}$  are independently selected from:

- 15 a) hydrogen,  
 b)  $C_1-C_6$  alkyl unsubstituted or substituted by alkenyl,  $R^{10}O-$ ,  
 $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $N_3$ ,  $(R^{10})_2N-C(NR^{10})-$ ,  
 $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ ,  
 20 c) aryl, heterocycle, cycloalkyl, alkenyl,  $R^{10}O-$ ,  
 $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $NO_2$ ,  
 $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  
 $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ , and  
 d)  $C_1-C_6$  alkyl substituted with an unsubstituted or  
 substituted group selected from aryl, heterocyclic and  
 $C_3-C_{10}$  cycloalkyl;

25  $R^4$  and  $R^5$  are independently selected from:

- a) hydrogen, and  
 b)



30  $R^7$  is independently selected from:

- a) hydrogen,

- 98 -

- b) aryl, heterocycle, cycloalkyl, alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $NO_2$ ,  $R^{10}_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ , and
- 5 c)  $C_1$ - $C_6$  alkyl unsubstituted or substituted by aryl, heterocycle, cycloalkyl, alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NH-$ , CN,  $H_2N-C(NH)-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{10}OC(O)NH-$ ;
- 10  $R^8$  is selected from:
- a) hydrogen,
- b) alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $NO_2$ ,  $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ , and
- 15 c)  $C_1$ - $C_6$  alkyl unsubstituted or substituted by perfluoroalkyl, F, Cl, Br,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ ;
- 20

$R^{10}$  is independently selected from hydrogen,  $C_1$ - $C_6$  alkyl, benzyl and aryl;

25  $R^{11}$  is independently selected from  $C_1$ - $C_6$  alkyl and aryl;

$A^1$  and  $A^2$  are independently selected from: a bond,  $-CH=CH-$ ,  $-C\equiv C-$ ,  $-C(O)-$ ,  $-C(O)NR^{10}-$ ,  $-NR^{10}C(O)-$ , O,  $-N(R^{10})-$ ,  $-S(O)_2N(R^{10})-$ ,  $-N(R^{10})S(O)_2-$ , or  $S(O)_m$ ;

30

V is selected from:

- a) hydrogen,
- b) heterocycle,
- c) aryl,



- 99 -

d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and

e) C<sub>2</sub>-C<sub>20</sub> alkenyl,

provided that V is not hydrogen if A<sup>1</sup> is S(O)<sub>m</sub> and V is not hydrogen if

5 A<sup>1</sup> is a bond, n is 0 and A<sup>2</sup> is S(O)<sub>m</sub>;

W is a heterocycle;

Z is independently H<sub>2</sub> or O;

10

m is 0, 1 or 2;

n is 0, 1, 2, 3 or 4;

p is 0, 1, 2, 3 or 4;

q is 0, 1 or 2;

15 r is 0 to 5, provided that r is 0 when V is hydrogen;


s is 4 or 5; and

u is 0 or 1;

or a pharmaceutically acceptable salt thereof.

20

4. A prodrug of the compound of Claim 3 having the Formula IV:



25

wherein:

R<sup>1a</sup> and R<sup>1b</sup> are independently selected from:

a) hydrogen,

- 100 -

- b) aryl, heterocycle, cycloalkyl, alkenyl, alkynyl,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $NO_2$ ,  $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ ,
- 5 c)  $C_1-C_6$  alkyl unsubstituted or substituted by aryl, heterocyclic, cycloalkyl, alkenyl, alkynyl,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ ;


10

$R^{2a}$  and  $R^{2b}$  are independently selected from:

- a) hydrogen,
- b)  $C_1-C_6$  alkyl unsubstituted or substituted by alkenyl,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $N_3$ ,  $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ ,
- 15 c) aryl, heterocycle, cycloalkyl, alkenyl,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ , CN,  $NO_2$ ,  $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ , and
- 20 d)  $C_1-C_6$  alkyl substituted with an unsubstituted or substituted group selected from aryl, heterocyclic and  $C_3-C_{10}$  cycloalkyl;

$R^4$  and  $R^5$  are independently selected from:

- 25 a) hydrogen, and
- b)



$R^7$  is independently selected from:

- 30 a) hydrogen,
- b) aryl, heterocycle, cycloalkyl, alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,

- 101 -

- $R^{10}C(O)NR^{10}-$ ,  $CN$ ,  $NO_2$ ,  $R^{10}_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  
 $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ , and  
 c)  $C_1$ - $C_6$  alkyl unsubstituted or substituted by aryl,  
 heterocycle, cycloalkyl, alkenyl, alkynyl, perfluoroalkyl, F,  
 5 Cl, Br,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NH-$ ,  $CN$ ,  $H_2N-$   
 $C(NH)-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ , or  
 $R^{10}OC(O)NH-$ ;

$R^8$  is selected from:

- 10 a) hydrogen,  
 b) alkenyl, alkynyl, perfluoroalkyl, F, Cl, Br,  $R^{10}O-$ ,  
 $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ ,  $CN$ ,  $NO_2$ ,  
 $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  
 $-N(R^{10})_2$ , or  $R^{11}OC(O)NR^{10}-$ , and  
 15 c)  $C_1$ - $C_6$  alkyl unsubstituted or substituted by perfluoroalkyl,  
 F, Cl, Br,  $R^{10}O-$ ,  $R^{11}S(O)_m-$ ,  $R^{10}C(O)NR^{10}-$ ,  $CN$ ,  
 $(R^{10})_2N-C(NR^{10})-$ ,  $R^{10}C(O)-$ ,  $R^{10}OC(O)-$ ,  $N_3$ ,  $-N(R^{10})_2$ ,  
 or  $R^{11}OC(O)NR^{10}-$ ;

- 20  $R^{10}$  is independently selected from hydrogen,  $C_1$ - $C_6$  alkyl, benzyl and aryl;

$R^{11}$  is independently selected from  $C_1$ - $C_6$  alkyl and aryl;

- 25  $A^1$  and  $A^2$  are independently selected from: a bond,  $-CH=CH-$ ,  $-C\equiv C-$ ,  
 $-C(O)-$ ,  $-C(O)NR^{10}-$ ,  $-NR^{10}C(O)-$ , O,  $-N(R^{10})-$ ,  
 $-S(O)_2N(R^{10})-$ ,  $-N(R^{10})S(O)_2-$ , or  $S(O)_m$ ;

V is selected from:

- 30 a) hydrogen,  
 b) heterocycle,  
 c) aryl,  
 d)  $C_1$ - $C_{20}$  alkyl wherein from 0 to 4 carbon atoms are replaced  
 with a heteroatom selected from O, S, and N, and

- 102 -

e) C<sub>2</sub>-C<sub>20</sub> alkenyl,  
provided that V is not hydrogen if A<sup>1</sup> is S(O)<sub>m</sub> and V is not hydrogen if  
A<sup>1</sup> is a bond, n is 0 and A<sup>2</sup> is S(O)<sub>m</sub>;


5 W is a heterocycle;

Z is independently H<sub>2</sub> or O;

m is 0, 1 or 2;  
10 n is 0, 1, 2, 3 or 4;  
p is 0, 1, 2, 3 or 4;  
q is 0, 1 or 2;  
r is 0 to 5, provided that r is 0 when V is hydrogen;  
s is 4 or 5; and  
15 u is 0 or 1;

or a pharmaceutically acceptable salt thereof.

5. The compound according to Claim 1 of the formula  
20 Ia:



wherein:

25 R<sup>1a</sup> is independently selected from: hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl;

R<sup>1b</sup> is independently selected from:

- a) hydrogen,
- b) aryl, heterocycle, cycloalkyl, R<sup>10</sup>O-, -N(R<sup>10</sup>)<sub>2</sub> or alkenyl,

- 103 -

- c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl, heterocycle, cycloalkyl, alkenyl, R<sup>10</sup>O-, or -N(R<sup>10</sup>)<sub>2</sub>;

R<sub>2a</sub> is selected from:

- 5        a) hydrogen,  
           b) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by alkenyl, R<sup>10</sup>O-,  
              R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, N<sub>3</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-,  
              R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-,  
 10        c) aryl, heterocycle, cycloalkyl, alkenyl, R<sup>10</sup>O-,  
              R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>,  
              (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>,  
              -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and  
           d) C<sub>1</sub>-C<sub>6</sub> alkyl substituted with an unsubstituted or  
              substituted group selected from aryl, heterocyclic and  
 15        C<sub>3</sub>- C<sub>10</sub> cycloalkyl;

R<sub>2b</sub> is hydrogen;


R<sub>3a</sub> and R<sub>3b</sub> are independently selected from:

- 20        a) a side chain of a naturally occurring amino acid,  
           b) an oxidized form of a side chain of a naturally occurring  
              amino acid which is:  
              i) methionine sulfoxide, or  
              ii) methionine sulfone,  
 25        c) substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>2</sub>-C<sub>10</sub> alkenyl,  
              C<sub>3</sub>-C<sub>10</sub> cycloalkyl, aryl or heterocyclic group,  
              wherein the substituent is selected from F, Cl, Br,  
              NO<sub>2</sub>, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN,  
              (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>,  
 30        -N(R<sup>10</sup>)<sub>2</sub>, R<sup>11</sup>OC(O)NR<sup>10</sup>- and C<sub>1</sub>-C<sub>20</sub> alkyl, and  
           d) C<sub>1</sub>-C<sub>6</sub> alkyl substituted with an unsubstituted or  
              substituted group selected from aryl, heterocycle and C<sub>3</sub>-  
              C<sub>10</sub> cycloalkyl;

- 104 -

R<sup>4</sup> and R<sup>5</sup> are independently selected from:

- a) hydrogen, and
- b)



5

R<sup>7</sup> is independently selected from:

- a) hydrogen,
- b) C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and
- c) C<sub>1</sub>-C<sub>6</sub> alkyl substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>10</sup>O-, R<sup>10</sup>C(O)NR<sup>10</sup>-, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-;

15

R<sup>8</sup> is selected from:

- a) hydrogen,
- b) C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and
- c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-;

25

R<sup>10</sup> is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl and aryl;

30 R<sup>11</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

A<sup>1</sup> and A<sup>2</sup> are independently selected from: a bond, -CH=CH-, -C≡C-,

- 105 -

-C(O)-, -C(O)NR<sup>10</sup>-, O, -N(R<sup>10</sup>)-, or S(O)<sub>m</sub>;

V is selected from:

- a) hydrogen,
- 5      b) heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, isoquinolinyl, and thienyl,
- c) aryl,
- 10      d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
- e) C<sub>2</sub>-C<sub>20</sub> alkenyl, and

provided that V is not hydrogen if A<sup>1</sup> is S(O)<sub>m</sub> and V is not hydrogen if A<sup>1</sup> is a bond, n is 0 and A<sup>2</sup> is S(O)<sub>m</sub>;

- 15      W is a heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, or isoquinolinyl;

Z is independently H<sub>2</sub> or O;

20


- m is            0, 1 or 2;
- n is            0, 1, 2, 3 or 4;
- p is            0, 1, 2, 3 or 4;
- r is            0 to 5, provided that r is 0 when V is hydrogen; and
- 25      u is            0 or 1;

or a pharmaceutically acceptable salt thereof.

6.      The compound according to Claim 2 of the formula

30      IIa:

- 106 -



R1a is independently selected from: hydrogen or C1-C6 alkyl;

5 R1b is independently selected from:

- a) hydrogen,
- b) aryl, heterocycle, cycloalkyl, R10O-, -N(R10)2 or alkenyl,
- c) C1-C6 alkyl unsubstituted or substituted by aryl, heterocycle, cycloalkyl, alkenyl, R10O-, or -N(R10)2;

10

R2a is selected from:

- a) hydrogen,
- b) C1-C6 alkyl unsubstituted or substituted by alkenyl, R10O-, R11S(O)m-, R10C(O)NR10-, CN, N3, (R10)2N-C(NR10)-, R10C(O)-, R10OC(O)-, -N(R10)2, or R11OC(O)NR10-,
- c) aryl, heterocycle, cycloalkyl, alkenyl, R10O-, R11S(O)m-, R10C(O)NR10-, CN, NO2, (R10)2N-C(NR10)-, R10C(O)-, R10OC(O)-, N3, -N(R10)2, or R11OC(O)NR10-, and
- d) C1-C6 alkyl substituted with an unsubstituted or substituted group selected from aryl, heterocyclic and C3- C10 cycloalkyl;

15

20

R2b is hydrogen;

25

R3a and R3b are independently selected from:

- a) a side chain of a naturally occurring amino acid,
- b) an oxidized form of a side chain of a naturally occurring amino acid which is:




- 107 -

- i) methionine sulfoxide, or  
 ii) methionine sulfone,  
 c) substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>2</sub>-C<sub>10</sub> alkenyl,  
 C<sub>3</sub>-C<sub>10</sub> cycloalkyl, aryl or heterocyclic group,  
 5 wherein the substituent is selected from F, Cl, Br,  
 NO<sub>2</sub>, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN,  
 (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>,  
 -N(R<sup>10</sup>)<sub>2</sub>, R<sup>11</sup>OC(O)NR<sup>10</sup>- and C<sub>1</sub>-C<sub>20</sub> alkyl, and  
 d) C<sub>1</sub>-C<sub>6</sub> alkyl substituted with an unsubstituted or  
 10 substituted group selected from aryl, heterocycle and C<sub>3</sub>-  
 C<sub>10</sub> cycloalkyl;

R<sup>4</sup> and R<sup>5</sup> are independently selected from:

- a) hydrogen, and  
 15 b)




R<sup>6</sup> is

- a) substituted or unsubstituted C<sub>1</sub>-C<sub>8</sub> alkyl or substituted or  
 20 unsubstituted C<sub>5</sub>-C<sub>8</sub> cycloalkyl, wherein the substituent on  
 the alkyl is selected from:  
 1) aryl,  
 2) heterocycle,  
 3) -N(R<sup>11</sup>)<sub>2</sub>,  
 25 4) -OR<sup>10</sup>, or

- 108 -

b)

R<sup>7</sup> is independently selected from:

- 5      a) hydrogen,  
       b) C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and  
 10      c) C<sub>1</sub>-C<sub>6</sub> alkyl substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>10</sup>O-, R<sup>10</sup>C(O)NR<sup>10</sup>-, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-;

R<sup>8</sup> is selected from:

- 15      a) hydrogen,  
       b) C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and  
 20      c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-;

25      R<sup>10</sup> is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl and aryl;

R<sup>11</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

30      R<sup>12</sup> is independently selected from hydrogen and C<sub>1</sub>-C<sub>6</sub> alkyl;

R<sup>13</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl;

- 109 -

A<sup>1</sup> and A<sup>2</sup> are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR<sup>10</sup>-, O, -N(R<sup>10</sup>)-, or S(O)<sub>m</sub>;

V is selected from:

- 5           a) hydrogen,  
          b) heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, isoquinolinyl, and thienyl,  
          c) aryl,  
10          d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and  
          e) C<sub>2</sub>-C<sub>20</sub> alkenyl, and

provided that V is not hydrogen if A<sup>1</sup> is S(O)<sub>m</sub> and V is not hydrogen if A<sup>1</sup> is a bond, n is 0 and A<sup>2</sup> is S(O)<sub>m</sub>;

15

W is a heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, or isoquinolinyl;


20   Z is independently H<sub>2</sub> or O;

- m is           0, 1 or 2;  
n is           0, 1, 2, 3 or 4;  
p is           0, 1, 2, 3 or 4;  
25   r is           0 to 5, provided that r is 0 when V is hydrogen; and  
u is           0 or 1;

or a pharmaceutically acceptable salt thereof.

30                   7.    The compound according to Claim 3 of the formula IIIa:

- 110 -



wherein:

R<sup>1a</sup> is independently selected from: hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl;

5 R<sup>1b</sup> is independently selected from:

- a) hydrogen,
- b) aryl, heterocycle, cycloalkyl, R<sup>10</sup>O-, -N(R<sup>10</sup>)<sub>2</sub> or alkenyl,
- c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl, heterocycle, cycloalkyl, alkenyl, R<sup>10</sup>O-, or -N(R<sup>10</sup>)<sub>2</sub>;

10

R<sup>2a</sup> is selected from:

- a) hydrogen,
- b) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by alkenyl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, N<sub>3</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-,
- 15 c) aryl, heterocycle, cycloalkyl, alkenyl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and
- 20 d) C<sub>1</sub>-C<sub>6</sub> alkyl substituted with an unsubstituted or substituted group selected from aryl, heterocyclic and C<sub>3</sub>- C<sub>10</sub> cycloalkyl;

R<sup>2b</sup> is hydrogen;


25

R<sup>4</sup> and R<sup>5</sup> are independently selected from:

- a) hydrogen, and

- 111 -

b)

R<sup>7</sup> is independently selected from:

- 5        a) hydrogen,
- b) C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and
- 10       c) C<sub>1</sub>-C<sub>6</sub> alkyl substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>10</sup>O-, R<sup>10</sup>C(O)NR<sup>10</sup>-, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-;

R<sup>8</sup> is selected from:

- 15       a) hydrogen,
- b) C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and
- 20       c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-;

- 25       R<sup>10</sup> is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl and aryl;

R<sup>11</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

- 30       A<sup>1</sup> and A<sup>2</sup> are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR<sup>10</sup>-, O, -N(R<sup>10</sup>)-, or S(O)<sub>m</sub>;

- 112 -

V is selected from:

- a) hydrogen,
  - b) heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, isoquinolinyl, and thienyl,
  - c) aryl,
  - d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
  - e) C<sub>2</sub>-C<sub>20</sub> alkenyl, and
- 10 provided that V is not hydrogen if A<sup>1</sup> is S(O)<sub>m</sub> and V is not hydrogen if A<sup>1</sup> is a bond, n is 0 and A<sup>2</sup> is S(O)<sub>m</sub>;

15 W is a heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, or isoquinolinyl;

Z is independently H<sub>2</sub> or O;


- m is 0, 1 or 2;
- 20 n is 0, 1, 2, 3 or 4;
- p is 0, 1, 2, 3 or 4;
- q is 0, 1 or 2;
- r is 0 to 5, provided that r is 0 when V is hydrogen; and
- u is 0 or 1;

25

or a pharmaceutically acceptable salt thereof.

8. The compound according to Claim 4 of the formula IVa:

- 113 -



wherein:

R<sup>1a</sup> is independently selected from: hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl;

5

R<sup>1b</sup> is independently selected from:

- a) hydrogen,
  - b) aryl, heterocycle, cycloalkyl, R<sup>10</sup>O-, -N(R<sup>10</sup>)<sub>2</sub> or alkenyl,
  - c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by aryl,
- 10 heterocycle, cycloalkyl, alkenyl, R<sup>10</sup>O-, or -N(R<sup>10</sup>)<sub>2</sub>;

R<sup>2a</sup> is selected from:

- a) hydrogen,
  - b) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by alkenyl, R<sup>10</sup>O-,
- 15 R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, N<sub>3</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-,  
R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-,
- c) aryl, heterocycle, cycloalkyl, alkenyl, R<sup>10</sup>O-,
- 20 R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>,  
(R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-,  
N<sub>3</sub>, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and
- d) C<sub>1</sub>-C<sub>6</sub> alkyl substituted with an unsubstituted or
- substituted group selected from aryl, heterocyclic and  
C<sub>3</sub>- C<sub>10</sub> cycloalkyl;


25 R<sup>2b</sup> is hydrogen;

R<sup>4</sup> and R<sup>5</sup> are independently selected from:

- a) hydrogen, and

- 114 -

b)

R<sup>7</sup> is independently selected from:

- 5        a) hydrogen,  
        b) C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and  
 10       c) C<sub>1</sub>-C<sub>6</sub> alkyl substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, R<sup>10</sup>O-, R<sup>10</sup>C(O)NR<sup>10</sup>-, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-;

R<sup>8</sup> is selected from:

- 15       a) hydrogen,  
        b) C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, NO<sub>2</sub>, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-, and  
 20       c) C<sub>1</sub>-C<sub>6</sub> alkyl unsubstituted or substituted by C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, F, Cl, R<sup>10</sup>O-, R<sup>11</sup>S(O)<sub>m</sub>-, R<sup>10</sup>C(O)NR<sup>10</sup>-, CN, (R<sup>10</sup>)<sub>2</sub>N-C(NR<sup>10</sup>)-, R<sup>10</sup>C(O)-, R<sup>10</sup>OC(O)-, -N(R<sup>10</sup>)<sub>2</sub>, or R<sup>11</sup>OC(O)NR<sup>10</sup>-;

- 25       R<sup>10</sup> is independently selected from hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl and aryl;

R<sup>11</sup> is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl and aryl;

- 30       A<sup>1</sup> and A<sup>2</sup> are independently selected from: a bond, -CH=CH-, -C≡C-, -C(O)-, -C(O)NR<sup>10</sup>-, O, -N(R<sup>10</sup>)-, or S(O)<sub>m</sub>;



- 115 -

V is selected from:

- a) hydrogen,
  - b) heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, isoquinolinyl, and thienyl,
  - c) aryl,
  - d) C<sub>1</sub>-C<sub>20</sub> alkyl wherein from 0 to 4 carbon atoms are replaced with a heteroatom selected from O, S, and N, and
  - e) C<sub>2</sub>-C<sub>20</sub> alkenyl, and
- provided that V is not hydrogen if A<sup>1</sup> is S(O)<sub>m</sub> and V is not hydrogen if A<sup>1</sup> is a bond, n is 0 and A<sup>2</sup> is S(O)<sub>m</sub>;

W is a heterocycle selected from pyrrolidinyl, imidazolyl, pyridinyl, thiazolyl, pyridonyl, 2-oxopiperidinyl, indolyl, quinolinyl, or isoquinolinyl;

Z is independently H<sub>2</sub> or O;

- m is 0, 1 or 2;
- n is 0, 1, 2, 3 or 4;
- p is 0, 1, 2, 3 or 4;
- q is 0, 1 or 2;
- r is 0 to 5, provided that r is 0 when V is hydrogen; and
- u is 0 or 1;

or a pharmaceutically acceptable salt thereof.

9. A compound which inhibits farnesyl-protein transferase which is:

N-(1(S)-carboxy-3-methylthiopropyl)-3-(4-imidazolylmethyl)-aminomethylbenzamide

- 116 -

N-(1(S)-carboxy-3-methylthiopropyl)-3-[N,N-*bis*-(4-imidazole-methyl)aminomethyl]benzamide

5 N-(1(S)-carboxy-3-methylthiopropyl) 3-[(4-imidazolylmethyl) -N-methylaminomethyl]benzamide

N-(1(S)-carboxy-3-methylthiopropyl) 4-[(4-imidazolylmethyl)-amino]benzamide

10 N-(1(S)-carboxy-3-methylthiopropyl)-3-[(4-imidazolylmethyl) amino] benzamide

N-(1(S)-Carboxy-3-methylthiopropyl)-3-[(4-imidazolylpropyl) amino] benzamide

15 N-(1(S)-Carboxy-3-methylthiopropyl)-3-[N-(4-imidazolylmethyl)-N-(4-nitrobenzyl)aminomethyl]benzamide

20 N-(1(S)-carboxy-3-methylthiopropyl)-3-[N,N-*bis*-(4-nitrophenylmethyl)aminomethyl]benzamide or

N-(1(S)-carbomethoxy-3-methylthiopropyl)-3-(4-imidazolylmethyl)-aminomethylbenzamide

25 N-(1(S)-carbomethoxy-3-methylthiopropyl)-3-[N,N-*bis*-(4-imidazole-methyl)aminomethyl]benzamide

N-(1(S)-carbomethoxy-3-methylthiopropyl) 3-[(4-imidazolylmethyl) -N-methylaminomethyl]benzamide

30 N-(1(S)-carbomethoxy-3-methylthiopropyl) 4-[(4-imidazolylmethyl)-amino]benzamide

- 117 -

N-(1(S)-carbomethoxy-3-methylthiopropyl)-3-[(4-imidazolylmethyl) amino] benzamide

5 N-(1(S)-carbomethoxy-3-methylthiopropyl)-3-[(4-imidazolylpropyl) amino]benzamide


N-(1(S)-Carbomethoxy-3-methylthiopropyl)-3-[N-(4-imidazolylmethyl)-N-(4-nitrobenzyl)aminomethyl]benzamide

10 N-(1(S)-carbomethoxy-3-methylthiopropyl)-3-[N,N-bis-(4-nitrophenylmethyl)aminomethyl]benzamide

or a pharmaceutically acceptable salt thereof.

15 10. A compound according to Claim 9 which inhibits farnesyl-protein transferase which is:

N-(1(S)-carboxy-3-methylthiopropyl)-3-[N,N-bis-(4-nitrophenylmethyl)]aminomethylbenzamide




20

or a pharmaceutically acceptable salt thereof.

11. A compound according to Claim 9 which inhibits farnesyl-protein transferase which is:


25 N-(1(S)-carbomethoxy-3-methylthiopropyl)-3-[N,N-bis-(4-nitrophenylmethyl)aminomethyl]benzamide

- 118 -



or a pharmaceutically acceptable salt thereof.


12. A compound according to Claim 9 which inhibits  
 5 farnesyl-protein transferase which is:  
 N-(1(S)-carboxy-3-methylthiopropyl)-3-[N,N-bis(4-imidazole-  
 methyl)aminomethyl] benzamide



- 10 or a pharmaceutically acceptable salt thereof.


13. A compound according to Claim 9 which inhibits  
 farnesyl-protein transferase which is:  
 N-(1(S)-carboxymethoxy-3-methylthiopropyl)-3-[N,N-bis(4-imidazole-  
 15 methyl)aminomethyl] benzamide

- 119 -



or a pharmaceutically acceptable salt thereof.


- 5                    14. A compound according to Claim 9 which inhibits farnesyl-protein transferase which is:  
N-(1(S)-Carboxy-3-methylthiopropyl)-3-[N-(4-imidazolylmethyl)-N-(4-nitrobenzyl)aminomethyl]benzamide



- 10 or a pharmaceutically acceptable salt thereof.

- 15                    15. A compound according to Claim 9 which inhibits farnesyl-protein transferase which is:  
N-(1(S)-Carbomethoxy-3-methylthiopropyl)-3-[N-(4-imidazolylmethyl)-N-(4-nitrobenzyl)aminomethyl]benzamide

- 120 -



or a pharmaceutically acceptable salt thereof.

5

16. A pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of Claim 2.

10

17. A pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of Claim 4.

15

18. A method for inhibiting farnesylation of Ras protein which comprises administering to a mammal in need thereof a therapeutically effective amount of the composition of Claim 9.

20

19. A method for inhibiting farnesylation of Ras protein which comprises administering to a mammal in need thereof a therapeutically effective amount of the composition of Claim 16.

25

20. A method for inhibiting farnesylation of Ras protein which comprises administering to a mammal in need thereof a therapeutically effective amount of the composition of Claim 17.

- 121 -

21. A method for inhibiting farnesylation of Ras protein which comprises administering to a mammal in need thereof a therapeutically effective amount of the composition of Claim 18.

5 22. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 16.

10 23. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 17.

15 24. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 18.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US96/03980

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 31/415; C07C 321/02; C07D 233/56  
US CL : Please See Extra Sheet.  
According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : Please See Extra Sheet.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A, E	US, A, 5,514,677 A (DAVIDSON et al.) 07 May 1996, see entire document.	1-24



Further documents are listed in the continuation of Box C.



See patent family annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be part of particular relevance
- \*E\* earlier document published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\*

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\*

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\*

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*G\*

document member of the same patent family

Date of the actual completion of the international search

17 JUNE 1996

Date of mailing of the international search report

11 JUL 1996

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JOSEPH K. MCKANE

Telephone No. (703) 308-1235



# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US96/03980

## A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

514/397, 399, 400, 561, 562, 563, 564, 570, 617, 618, 619; 548/312.7, 338.1; 562/426, 435, 443, 444, 470; 564/164

## B. FIELDS SEARCHED

Minimum documentation searched

Classification System: U.S.

A61K 31/415; C07C 321/02; C07D 233/56

514/397, 399, 400, 561, 562, 563, 564, 570, 617, 618, 619; 548/312.7, 338.1; 562/426, 435, 443, 444, 470; 564/164